**ACTIVITY COEFFICIENTS OF GLYCINE,
DL-SERINE AND DL-VALINE IN AQUEOUS
SOLUTIONS CONTAINING NITRATES AT 298.15 K**

by
Chongxia Gao

Department of Chemical Engineering
McGill University, Montreal, Canada

A thesis submitted to the Faculty of Graduate Studies and Research in partial
fulfillment of the requirements of the degree of Master of Engineering

© Chongxia Gao March, 2000

**National Library
of Canada**

**Acquisitions and
Bibliographic Services**

**395 Wellington Street
Ottawa ON K1A 0N4
Canada**

**Bibliothèque nationale
du Canada**

**Acquisitions et
services bibliographiques**

**395, rue Wellington
Ottawa ON K1A 0N4
Canada**

Your file *Votre référence*

Our file *Notre référence*

The author has granted a non-exclusive licence allowing the National Library of Canada to reproduce, loan, distribute or sell copies of this thesis in microform, paper or electronic formats.

L'auteur a accordé une licence non exclusive permettant à la Bibliothèque nationale du Canada de reproduire, prêter, distribuer ou vendre des copies de cette thèse sous la forme de microfiche/film, de reproduction sur papier ou sur format électronique.

The author retains ownership of the copyright in this thesis. Neither the thesis nor substantial extracts from it may be printed or otherwise reproduced without the author's permission.

L'auteur conserve la propriété du droit d'auteur qui protège cette thèse. Ni la thèse ni des extraits substantiels de celle-ci ne doivent être imprimés ou autrement reproduits sans son autorisation.

0-612-64220-8

Canada

To My Family

ABSTRACT

Both an electrochemical and an isopiestic methods were used to measure the mean activity coefficients of amino acids in water-electrolyte-amino acid systems. The electrochemical method used an electrochemical cell with an anion and a cation ion selective electrodes, each measured versus a double junction reference electrode. The activity coefficients of the electrolyte were obtained from the e.m.f measurements for four systems at 298.15K. The amino acids were glycine, DL-serine and DL-valine and the electrolyte were KNO_3 and NaNO_3 . The isopiestic method was applied to measure the activity of water for the system water + DL-serine + KNO_3 at 298.15K and the results were compared with those from electrochemical method. The activity coefficients of the amino acids in the ternary systems were obtained from those of the electrolyte measured by the electrochemical method or from those of the water measured by isopiestic method. according to the cross differential relation.

Two models were used to correlate the measured data. First the excess Gibbs free energy model proposed by Khoshkbarchi and Vera (1996a) with the NRTL equation was used to correlate the activity coefficient data. A simple model derived from a modification of a more complex model based on the perturbation theory (Khoshkbarchi and Vera, 1996c) was proposed. The model was applied to correlate the activity coefficients and solubilities of amino acids in binary aqueous solutions and the activity coefficients in ternary systems with satisfactory results.

RÉSUMÉ

Les méthodes électrochimique et isopiétique sont utilisées pour mesurer les coefficients d'activité des acides aminés dans une solution eau - électrolyte - acide aminé. La méthode électrochimique consiste en une cellule électrochimique formée d'une anode et d'une cathode, électrode sélective des ions chacun étant mesurée versus d'une électrode de référence à double jonction. Les coefficients d'activité de l'électrolyte sont obtenus grâce à la force électromotrice (e.m.f) pour quatre systèmes à 298.15K. Les acides aminés étudiés étaient: glycine, DL-serine et DL-valine et les électrolytes étaient KNO_3 et NaNO_3 . La méthode isopiétique est employée pour mesurer l'activité de l'eau dans le système eau - DL-serine - KNO_3 à 298.15K et les valeurs sont comparées avec la méthode électrochimique. Les coefficients d'activité des acides aminés dans le système ternaire proviennent de l'électrolyte mesuré par la méthode électrochimique, ou dans le cas de la méthode isopiétique, de l'eau mesurée, conformément à la relation de différentiation.

Les données recueillies sont corrélées par deux modèles. En premier lieu le modèle, basé sur l'excès d'énergie libre de Gibbs, proposé par Khoshkbarchi et Vera (1996 a) avec l'équation NRTL, a été employé pour corréler les résultats du coefficient d'activité. Un modèle simple dérivé d'un modèle plus complexe fondé sur la théorie de perturbation (Khoshkbarchi et Vera, 1996 c) est développé. Le modèle est utilisé avec succès pour corréler les coefficients d'activité et les solubilités des acides aminés dans des solutions aqueuses binaires et les coefficients d'activité dans des solutions ternaires.

ACKNOWLEDGMENTS

I am deeply indebted to Professor Juan H. Vera, my supervisor, not only for giving me the opportunity to work in his group, his support and his guidance throughout the course of this research, but also for his enthusiasm, encouragement, patience and kindness. I have learned from him more than thermodynamics and research techniques.

I am grateful to the other Professors in our research group, Prof. M. E. Weber and Prof. J. Simandl for their interests, valuable comments and suggestions.

Thanks also go to many Ph. D. and Master students in this group and the department: R. Dumortier, M. Husein, A. Merchant, H. Niu, K. Pehrson, A. Pradhan, Y.-O. Shin, V. Taghikhani, E. Wong. Their help, encouragement and enthusiasm made my stay in this group an interesting and joyful experience.

Finally, I want to express my gratitude to the Chemical Engineering Department. Many people in the department gave me a lot of help during my research work.

TABLE OF CONTENTS

	Page
ABSTRACT	i
RESUME.....	ii
ACKNOWLEDGMENTS.....	iii
TABLE OF CONTENTS	iv
LIST OF FIGURES	vi
LIST OF TABLES.....	vii
LIST OF SYMBOLS	viii
 CHAPTER 1: INTRODUCTION	 1
1.1 General Remarks	1
1.2 Chemical Structure of Amino Acids	2
1.3 Experimental Measurement of Activity Coefficients	3
1.4 Modeling of Activity Coefficients	6
1.5 Objectives of Present Work	7
 CHAPTER 2: ELECTROCHEMICAL MEASUREMENT OF ACTIVITY COEFFICIENTS OF AMINO ACIDS IN AQUEOUS ELECTROLYTE SOLUTIONS	 10
2.1 Theoretical Basis of the Measurements	10
2.2 Experimental Set Up, Materials and Methods for Measuring the Activity Coefficients of Electrolytes in Aqueous Solutions	12
2.3 Experimental Results and Discussions	15
 CHAPTER 3: MEASUREMENT OF ACTIVITY COEFFICIENT USING ISOPIESTIC METHOD	 26

3.1	Motivation	26
3.2	Theoretical Background	26
3.3	Set Up and Measurements	28
3.4	Results and Discussions	30
3.5	Comparison with the Electrochemical Method	32
CHAPTER 4: MODELING		36
4.1	Gibbs Free Energy Model with NRTL Equation	36
4.2	A New Model Based on a Simplified Perturbation Theory	38
4.2.1	Theoretical Background.....	39
4.2.2	A New Model.....	41
4.3	Calculation Results.....	46
4.3.1	Binary systems.....	46
4.3.2	Ternary systems.....	48
4.4	Modeling the Solubility of Amino Acid in Water.....	51
CHAPTER 5: CONCLUSIONS, CONTRIBUTIONS TO KNOWLEDGE AND SUGGESTIONS FOR FUTURE WORK.....		55
5.1	Conclusions.....	55
5.2	Contributions to Knowledge.....	56
5.3	Suggestions for Future Work.....	56
REFERENCES.....		57
APPENDICES.....		62
A	ΔE data by Electrochemical Method.....	62
B	Derivation of Eq.(4-15).....	64
C	Derivation of the Perturbation Term for Mixtures.....	65
D	The Cross Differential Relation for the Perturbation Theory.....	66

LIST OF FIGURES

Figure 1.1 Structure of amino acids. (a) molecular structure of α -amino acid (b) structure of the zwitterion form.....	2
Figure 2.1 Schematic view of the set up used for measurement of the mean ionic activity coefficient of an electrolyte in aqueous solutions.....	13
Figure 2.2 Effect of concentrations of NaNO_3 and DL-serine, in aqueous solutions at 298.2 K, on the ratios $\gamma_{\pm}^{(2)}/\gamma_{\pm}^{(1)}$; $\gamma_A^{(2)}/\gamma_A^{(1)}$. Dots: experimental points; curves: equations (2-10), (2-13), respectively.....	20
Figure 2.3 Effect of concentrations of KNO_3 and DL-serine, in aqueous solutions at 298.2 K, on the ratios $\gamma_{\pm}^{(2)}/\gamma_{\pm}^{(1)}$; $\gamma_A^{(2)}/\gamma_A^{(1)}$. Dots: experimental points; curves: equations (2-10), (2-13), respectively.....	21
Figure 2.4 Effect of concentrations of NaNO_3 and DL-valine, in aqueous solutions at 298.2 K, on the ratios $\gamma_{\pm}^{(2)}/\gamma_{\pm}^{(1)}$; $\gamma_A^{(2)}/\gamma_A^{(1)}$. Dots: experimental points; curves: equations (2-10), (2-13), respectively.....	23
Figure 2.5 Re-plot of $\gamma_A^{(2)}/\gamma_A^{(1)}$ as a function of m_A for fixed values of m_S	24
Figure 2.6 Effect of concentrations of KNO_3 and glycine, in aqueous solutions at 298.2 K, on the ratios $\gamma_{\pm}^{(2)}/\gamma_{\pm}^{(1)}$; $\gamma_A^{(2)}/\gamma_A^{(1)}$. Dots: experimental points; curves: equations (2-10), (2-13), respectively.....	25
Figure 3.1 A view of isopiestic set up	30
Figure 3.2 Comparison of activity coefficients of DL-serine by two methods.....	35
Figure 3.3 Comparison of mean ionic activity coefficient by two methods.....	35
Figure 4.1 Correlation results for activity coefficients of amino acids in binary aqueous solutions.....	47
Figure 4.2 Correlation results by the simplified perturbation model for the mean activity coefficient of KNO_3 in ternary systems ($\text{KNO}_3 + \text{glycine} + \text{H}_2\text{O}$).....	50
Figure 4.3 Correlation results for solubility of amino acids in H_2O	54

LIST OF TABLES

Table 1.1: Systems for which activity coefficients of amino acids in aqueous electrolyte solutions have been reported.....	5
Table 1.2: Systems to be measured.....	7
Table 1.3 Chemical structure of the amino acids used in this work.....	8
Table 2.1 Experimental results for NaNO_3 + DL-serine system.....	16
Table 2.2 Experimental results for NaNO_3 + DL-valine system	16
Table 2.3 Experimental results for KNO_3 + DL-serine system	17
Table 2.4 Experimental results for KNO_3 + glycine system.....	17
Table 2.5 Pooled standard deviation of measured ratio $\gamma_{\pm}^{(2)}/\gamma_{\pm}^{(1)}$	18
Table 2.6 Coefficients of eqs.(2-10) and (2-13).....	18
Table 3.1 Correlation results by eq.(3-10).....	31
Table 3.2 Correlation results by eq.(3-11).....	32
Table 3.2 Measured data from isopiestic method.....	34
Table 3.3 Correlation results.....	34
Table 4.1 Values of pure component parameters of ions and electrolytes.....	45
Table 4.2 Calculation results for activity coefficients of amino acids.....	46
Table 4.3 Results for ternary systems by Gibbs free energy model.....	48
Table 4.4 Results for ternary systems by perturbation model.....	49
Table 4-5 Calculation results for solubility of amino acids.....	53
Table A.1 ΔE data for NaNO_3 + DL-serine + H_2O system.....	58
Table A.2 ΔE data for NaNO_3 + DL-valine + H_2O system.....	58
Table A.3 ΔE data for KNO_3 + DL-serine + H_2O system.....	59
Table A.4 ΔE data for KNO_3 + glycine + H_2O system.....	59

LIST OF SYMBOLS

a	Activity
A	Constant in eq.(3-10)
AAD	Average absolute deviation
b, c, d	Parameters in eq.(3-11)
B	Constant, 0.2
B_i	Parameters in eq.(3-10), $i=1,3$
c	Total number of components
C_i	Adjustable parameters in eqs.(2-10), (3-7), $i=0,6$
D	Dipole moment
e.m.f.	Electromotive force
E_+	Potential of the cation ion selective electrode
E_-	Potential of the anion ion selective electrode
E_+^{ISE}	Internal potential of the cation ion selective electrode
E_+^o	Standard state potential of the cation ion selective electrode
E_-^o	Standard state potential of the anion ion selective electrode
E_-^{ISE}	Internal potential of the anion ion selective electrode
E^J	Junction potential
E^{Ref}	Reference electrode potential
$\Delta E^{(1)}$	Potential difference in electrochemical cell in the presence of electrolyte and in the absence of amino acid
$\Delta E^{(2)}$	Potential difference in electrochemical cell in the presence of both electrolyte and amino acid
F	Faraday number
f	Fugacity
G	Gibbs free energy
G_{ij}	Binary interaction energy function in the NRTL model
$g(r)$	Radial distribution function
Δh	Enthalpy change
ISE	Ion selective electrode

k	Boltzmann constant ($1.381 \times 10^{-23} \text{ J K}^{-1}$)
m	Concentration in molality unit
m_A	Molality of amino acid
m_S	Molality of electrolyte
m_+	Molality of cation
m_-	Molality of anion
M	Molecular weight
n	Number of moles
n_D	Number of data points
N	Number of molecules
N_A	Avogadro number (6.02252×10^{23})
O.F.	Objective function
r.m.s.d.	Root mean square deviation
r	Intermolecular or interionic distance
R	Universal gas constant
Δs	Entropy change
S	Slope of electrode potential
S	Solubility of amino acid in water
s.e.e	Standard error of estimate
T	Absolute temperature
u	Intermolecular potential
V	Volume
x	Mole fraction
Z_+	Charge number of cation
Z_-	Charge number of anion

Greek letters

α	Parameter of the NRTL model (0.3)
$\bar{\alpha}$	Polarizability
η	Quantity defined by eq.(4-11)

$\delta^{(0)}, \delta^{(1)}$	Parameters in mixing rule eq.(4-27)
Δ	Quantity defined by eq.(3-6)
γ	Activity coefficient
γ_A	Activity coefficient of amino acid
γ_S	Activity coefficient of electrolyte
γ_+	Activity coefficient of cation
γ_-	Activity coefficient of anion
$\gamma_{\pm}$	Mean ionic activity coefficient of electrolyte
$\gamma_{\pm}^{(1)}$	Mean ionic activity coefficient of electrolyte in electrochemical cell in the absence of amino acid
$\gamma_{\pm}^{(2)}$	Mean ionic activity coefficient of electrolyte in electrochemical cell in the presence of amino acid
ε	Depth of potential well, molecular energy parameter
ε_0	Permittivity of the vacuum
ε_r	Relative dielectric constant
μ	Chemical potential
ν	Stoichiometric number of electrolyte, ($\nu_+ + \nu_-$)
ν_+	Stoichiometric number of cations
ν_-	Stoichiometric number of anions
ρ	Number density
σ	Molecular diameter, size parameter
τ_{ij}	Binary energy parameter between species i and j in the NRTL model
ϕ	Osmotic coefficient

Superscripts

0	Standard state
Cal	Calculated
E	Excess free energy
exp	Experimental

HS	Hard sphere
ISE	Ion selective electrode
K-V	K-V model
L	Liquid phase
L-J	Lennard-Jones
LR	Long range interactions term
Per	Perturbed term
Ref	Reference term
S	Solid phase
SR	Short range interactions term

Subscripts

A	Amino acid
e	Effective hard sphere diameter
i, j, k	Species i, j, k
W	Water
S	Electrolyte
+	Cation
-	Anion
$\pm$	Mean ionic

CHAPTER 1: INTRODUCTION

1.1 General Remarks

The industrial scale production of biochemicals is of potential interests for the chemical, pharmaceutical and food industries. The separation and concentration process of biomolecules is one of the economically important stages in their production processes. As amino acids are the simplest biomolecules and the building blocks of other biomolecules such as peptides and proteins, a knowledge of the thermodynamic properties of amino acids in aqueous electrolyte solutions is a first step towards the rational design of equilibrium-based separation processes. In addition, such information is useful for the understanding of the molecular interactions in these systems.

The experimental and theoretical research on the equilibrium properties of amino acid solutions have attracted attention in recent years. The chemical engineering thermodynamics research group at McGill University has published a series of articles on separation processes and thermodynamic properties of amino acid in aqueous electrolyte solutions. A new electrochemical method was developed for the measurement of activity coefficients of amino acids (Khoshkbarchi and Vera, 1996a,b). Data of activity coefficients and solubilities were published for various systems and some theoretical models were proposed for these properties (Khoshkbarchi and Vera, 1996a, b, c, d, e; Khoshkbarchi and Vera, 1997; Soto-Campos et al., 1997a,b, 1998; Pradhan and Vera, 1998). This work is a continuation of research work carried out by this research group.

1.2 Chemical Structure of Amino Acids

The physico-chemical behavior of amino acids in aqueous electrolyte solutions is the result of their special molecular structure. As shown in Fig.1 (a), an α -amino acid molecule is composed of three parts: a terminal carboxyl group (-COOH), an amino

Chapter 1 INTRODUCTION

group ($-\text{NH}_2$), and a hydrocarbon backbone ($-\text{R}$) with variable length, connected together by a central carbon atom (called α carbon, C_α). If the amino group is attached to the carbon next to the α carbon of the alkyl chain $-\text{R}$, the amino acid is called β -amino acid. Thus depending on the location of the amino group in the hydrocarbon backbone of the amino acid, they are classified as α , β , γ , δ , ϵ , etc. amino acids.

In aqueous solutions in the neutral pH range, amino acids form a zwitterion (zwitter = two) resulting from the transfer of a proton from the carboxyl group to the amino group. Therefore, an amino acid molecule possesses, at the same time, a negative and a positive charge and this form is called a zwitterion. The electric field surrounding these charged ions gives rise to important interactions. The structure of the zwitterion is shown in Fig. 1.1(b).

Figure 1.1 Structure of amino acids. (a) molecular structure of α -amino acid (b) structure of the zwitterion form.

The following reactions may take place when an amino acid is dissolved in water at different pH values:

Chapter 1 INTRODUCTION

In aqueous solution at neutral pH, the amino acid molecules are in their zwitterionic form and the concentration of anionic and cationic forms are negligible. At low pH, the amino molecules are in cationic form, NH_3^+ROOH , and at high pH they are in anionic form, NH_2RCOO^- .

1.3 Experimental Measurement of Activity Coefficients

There are two types of methods commonly used for measuring the activity coefficients of amino acid in aqueous electrolyte solutions. One is the electrochemical method and the other is the isopiestic method. Both methods are indirect, in the sense that the activity coefficient of the amino acid is obtained by measuring either the mean ionic activity coefficient of the electrolyte or the osmotic coefficient of the solution.

Electrochemical method

The electrochemical method has been widely used for measuring the activity coefficients of electrolytes in aqueous solutions. In some measurements of activity coefficients of amino acid-electrolyte-water systems using electrochemical method (Kelley and Lilley, 1978; Rodriguez-Raposo et al., 1994), only one ion selective electrode (usually a cation ion selective electrode) versus a reference electrode, was used to measure the mean ionic activity coefficients of electrolyte in the presence of another solute (i.e., amino acid). The drawbacks of this method have been discussed by Khoshkbarchi and Vera (1996a, b). A new method was proposed by Khoshkbarchi and Vera (1996a, b), in which two ISEs with the same reference electrode were used so as to eliminate the shortcomings of the single ISE method and obtain more reliable results. This latter method, using two IESs, will be used in this work. The major advantage of this method is that the measurement takes a very short time in comparison with the isopiestic method. Its disadvantage is that the measurements are limited by the range of applicability of the electrodes.

Chapter 1 INTRODUCTION

Isopiestic Method:

The isopiestic method was proposed more than a century ago and modified by Sinclair (1933). A detailed example of well designed apparatus was given by Scatchard et al. (1938). In 50's, many researchers used this methods (Robinson and Stokes, 1959). Two vessels were used in the measurements and the time needed for reaching equilibrium was very long. Recently, improved isopiestic methods were developed: Thiessen and Wilson (1987) used a three vessel apparatus; Ochs et al. (1990) used an apparatus with nine vessels, and a similar method was used by Lin et al. (1996) and by Park and Englezos (1998).

The major advantage of this method is that it can be used over the whole range of concentration. The main disadvantage is that it is time consuming, though an apparatus with multiple vessels may partially solve the problem.

In this work, an apparatus which is similar to the design used by Park and Englezos (1998) is used to measure the osmotic coefficients of aqueous electrolyte solutions containing amino acids. The results obtained are compared with those obtained by the electrochemical method for the system of DL-serine- KNO_3 - H_2O .

Systems studied

Table 1.1 lists the systems of amino acid-electrolyte -water for which activity coefficients have been reported according to our knowledge.

Table 1.1 Systems for which activity coefficients of amino acids in aqueous electrolyte solutions have been reported

Amino acid	Electrolyte	Method ^a	References ^b
glycine	NaCl	EC, IP	[1], [9], [13]
DL-alanine	NaCl	EC	[1]
DL-valine	NaCl	EC	[2]
DL-serine	NaCl	EC	[3]
glycine	KCl	EC	[4]
DL-alanine	KCl	EC	[5]
DL-valine	KCl	EC	[6]
DL-serine	KCl	EC	[3]
glycine	NaNO ₃	EC	[7]
DL-alanine	NaNO ₃	EC	[5]
DL-methionine	NaCl	EC	[7]
DL-threonine	NaCl	EC	[8]
DL-threonine	NaNO ₃	EC	[8]
β-alanine	NaCl	IP	[10]
γ-aminobutyric acid	NaCl	IP	[10]
γ-aminocaproic acid	NaCl	IP	[10]
L-α-alanine	NaCl	IP	[11]
glycine	CaCl ₂	EC	[12]
β-alanine	CaCl ₂	EC	[12]
γ-aminobutyric acid	CaCl ₂	EC	[12]
ε-aminocaproic acid	CaCl ₂	EC	[12]

a, EC: electrochemical method; IP: isopiestic method; b, see next page.

Chapter I INTRODUCTION

[1] Khoshkbarchi and Vera, 1996b; [2] Khoshkbarchi and Vera, 1996a; [3] Khoshkbarchi et al., J. 1997; [4] Bower and Robinson, 1965; [5] Soto -Campos et al., 1998; [6] Khoshkbarchi and Vera, 1996c; [7] Soto-Campos et al., 1997a; [8] Soto-Campos et al., 1997b; [9] Rodriguez-Raposo et al., 1994; [10] Schrier and Robinson, 1971; [11] Schrier and Robinson, 1974; [12] Briggs et al., 1974; [13] Phang and Steel, 1974.

Previous results show that the effect of salts on the activity of an amino acid is considerable. In some cases adding salt into the solution will increase the activity coefficient of the amino acid, while in other cases the activity coefficient of the amino acid will decrease as salt is added.

1.4 Modeling of Activity Coefficients

The modeling of activity coefficients is important since experimental methods only give data at discrete values of concentration and temperature. A reliable model can provide useful correlation and even prediction of results over a wide range of conditions based on a few experimental values.

Many theoretical or semi-empirical models have been proposed in literature for the systems containing electrolytes and amino acids. For example, Khoshkbarchi and Vera (1996a) proposed a Gibbs free energy model which used the Wilson or the NRTL equation for the short range interaction contributions. The same authors have also proposed a model based on the perturbation theory (Khoshkbarchi and Vera, 1996c,d). In this work, the Gibbs free energy model with the NRTL equation will be used to correlate the activity coefficient data for the ternary systems measured. In addition, a new simplified model based on the perturbation theory will be used to represent the activity coefficient of amino acids in both binary and ternary systems.

1.5 Objectives of the Present Work

The objectives of this work are the following:

(1a) To measure the activity coefficients of the electrolyte and calculate the activity coefficients of the amino acid at 298.15 K for the four systems listed in Table 1.2.

Table 1.2 Systems to be measured

	Amino acid	Electrolyte	Amino acid molality	Electrolyte molality	Method
1	glycine	KNO ₃	$0.05 < m_A < 2.50$	$0.1 < m_S < 0.7$	EC
2	DL-serine	KNO ₃	$0.05 < m_A < 0.40$	$0.1 < m_S < 0.7$	EC, IP
3	DL-valine	NaNO ₃	$0.05 < m_A < 0.50$	$0.1 < m_S < 1.0$	EC
4	DL-serine	NaNO ₃	$0.05 < m_A < 0.40$	$0.1 < m_S < 1.0$	EC

These ranges of molalities were chosen so to be within the concentration range in which the ISE electrodes are reliable and also to be below the solubility limit of the electrolyte and the amino acid in the system. The electrochemical method will be used in all cases.

(1b) To present a comparison of the results for DL-serine + KNO₃ system obtained from the electrochemical and isopiestic methods.

The chemical structure of the amino acids used in this work are given by Table 1.3.

Table 1.3 Chemical structure of the amino acids used in this work

Amino acid	Chemical structure
Glycine	
DL-serine	
DL-valine	

(2). To present a summary of the salt effects on the activities of amino acids in the systems measured and explain the effects based on the interactions between ions and zwitterions and their structures.

(3) To model the experimental data using the Gibbs free energy model proposed by Khoshkbarchi and Vera (1996a).

(4) To use a simplified model based on the perturbation theory to correlate activity coefficients of amino acids in both binary and ternary systems.

Chapter 1 INTRODUCTION

(5) To use the new activity coefficient model for the correlation of solubility of amino acids in water.

The rest of the thesis is organized as follows: Chapter 2 presents the principle of the measurement of activity coefficients using the electrochemical method, the experimental results obtained in this work and discussion. Chapter 3 presents the principle of the measurement of activity coefficients using the isopiestic method, the experimental results obtained in this work and discussion. Chapter 4 gives the derivation of a new model and results of the modeling using two approaches and discussion. Chapter 5 includes a general discussion of the work, the main contributions and suggestions for future work.

CHAPTER 2: ELECTROCHEMICAL MEASUREMENT OF ACTIVITY COEFFICIENTS OF AMINO ACIDS IN AQUEOUS ELECTROLYTE SOLUTIONS

2.1 Theoretical Basis of the Measurements

Among the various electrochemical methods, electrochemical cells with ion selective electrodes (ISE) are an efficient tool for the measurements of thermodynamic properties of electrolyte solutions (Haghtalab and Vera, 1991). The electrochemical method has proved to give reliable results for the activities of solutes in electrolyte solutions (Kelley and Lilley, 1978; Rodriguez-Raposo et al., 1994; Khoshkbarchi and Vera, 1996a, b).

The activity coefficients of the solute (amino acid) are obtained from measurements of the electrochemical potential (e.m.f) in electrochemical cells of the type:

It is also necessary to measure the e.m.f in an electrolyte aqueous solution without amino acid for calibration purposes. For the electrochemical cell:

the relation between e.m.f. of a cation selective electrode and activity coefficients of the cation is given by the Nerst equation:

$$E_{+} = E_{+}^{\circ} + \frac{RT}{z_{+}F} \ln(m_{+}\gamma_{+}) \quad (2-1)$$

Chapter 2. ELECTROCHEMICAL MEASUREMENT OF ACTIVITY COEFFICIENTS OF AMINO ACIDS...

$$E_- = E_-^0 - \frac{RT}{z_- F} \ln (m_- \gamma_-) \quad (2-2)$$

where R is the gas constant, T , temperature, z , the charge number, F the Faraday constant, m the molality of the ion, γ the activity coefficient, and E the e.m.f. The terms E_-^0 and E_-^0 are combinations of junction potential (E^J), reference electrode potential (E^{Ref}), and the ion selective electrode potential (E^{ISE}):

$$E_-^0 = E^{\text{Ref}} + E^J + E_-^{\text{ISE}} \quad (2-3)$$

$$E_-^0 = E^{\text{Ref}} + E^J + E_-^{\text{ISE}} \quad (2-4)$$

Combining eqs.(2-1), (2-2) and after rearrangement, we have

$$E_- - E_- = (E_-^0 - E_-^0) + S \ln(m_s \gamma_{\pm}) \quad (2-5)$$

where m_s is the molality of salt and the mean ionic activity coefficient of the electrolyte is defined as

$$\gamma_{\pm} = (\gamma_+^{\nu_+} \gamma_-^{\nu_-})^{1/(\nu_+ + \nu_-)} \quad (2-6)$$

In eq.(2-6), ν_+ and ν_- are the stoichiometric numbers of moles of cation and anion produced per mole of electrolyte. In eq.(2-5) the slope S is a function of temperature and should be calculated from linear fitting of the values of ΔE versus $\ln(m_s \gamma_{\pm})$ from a system for which activity coefficients of electrolyte are known. Applying eq.(2.5) to systems electrolyte + water (superscript 1) and amino acid + electrolyte + water (superscript 2) at the same molality of the electrolyte:

$$\Delta E^{(1)} = E^0 + S \ln(m_s \gamma_{\pm}^{(1)}) \quad (2-7)$$

$$\Delta E^{(2)} = E^0 + S \ln(m_s \gamma_{\pm}^{(2)}) \quad (2-8)$$

Chapter 2. ELECTROCHEMICAL MEASUREMENT OF ACTIVITY COEFFICIENTS OF AMINO ACIDS...

then

$$\ln\left(\frac{\gamma_{\pm}^{(2)}}{\gamma_{\pm}^{(1)}}\right) = \frac{\Delta E^{(2)} - \Delta E^{(1)}}{S} \quad (2-9)$$

Eq.(2.9) is used to calculate the mean ionic activity coefficient of the electrolyte in aqueous solutions containing an amino acid. The values of $\gamma_{\pm}$ in binary electrolyte + water systems are obtained from the literature.

In this work the differences of e.m.f., $\Delta E^{(1)}$ and $\Delta E^{(2)}$ in aqueous solutions have been measured using the method proposed by Khoshkbarchi and Vera (1996a) due to its advantages over the traditional method (see, e.g., Briggs et al., 1974). The electrochemical cell used is as follows:

Cation ISE | electrolyte (m_S) + solute (m_A) + water | reference electrode

Anion ISE | electrolyte (m_S) + solute (m_A) + water | reference electrode

The measured activity coefficients of the electrolytes are then used to evaluate the activity coefficients of the amino acid, as discussed below.

2.2 Experimental Set Up, Materials and Methods for Measuring the Activity Coefficients of Electrolytes in Aqueous Solutions

The experimental set up used in this work is illustrated in Figure 2.1. Basically it consists of a cation and an anion selective electrodes with a double junction reference electrode. A Ross sodium ion selective electrode, glass body model 84-11, and a double-junction electrode, model 13-620-46, were obtained from Orion (Boston, MA). A Potassium ion selective electrode, polymer body, model 93-19, was obtained from Radiometer (Kopenhagen, Denmark). A nitrate ion electrode, polymer body, model 9300BN was obtained from Orion. An Oakton pH/ISE meter (WD-35616-00, Singapore) with a resolution of ± 0.1 mV was used to monitor the e.m.f. measurements.

Chapter 2. ELECTROCHEMICAL MEASUREMENT OF ACTIVITY COEFFICIENTS OF AMINO ACIDS...

Figure 2.1 Schematic view of the set up used for measurement of the mean ionic activity coefficient of an electrolyte in aqueous solutions. 1: millivoltmeter; 2: reference electrode; 3: cation ISE; 4: cell; 5: magnetic stirrer; 6: magnet; 7: anion ISE; 8: thermometer; 9: thermostatic bath.

Chapter 2. ELECTROCHEMICAL MEASUREMENT OF ACTIVITY COEFFICIENTS OF AMINO ACIDS...

The chemicals used were obtained from A&C American Chemicals LTD (Montreal). The purity of salts used (NaNO_3 , KNO_3) was 99.9% and those of Glycine, DL- valine, and DL-serine were 98-99%.

The salts were oven-dried at 110 °C for 72 hours prior to use. The amino acids were used as received. The electrodes were stored and conditioned according to the manufacturer's instructions.

Experimental procedure

The aqueous solutions of electrolytes at desired molality were prepared and solid amino acids were then subsequently added. Deionized water with a conductivity less than 0.8 $\mu\text{S}/\text{cm}$ was used in all experiments. Water, electrolyte and amino acid were weighed. The compositions of the initial solutions were accurate to within ± 0.01 wt%. The volume of the solution was about 200 ml. The electrode was carefully prepared according to the instruction manual. The temperature was kept at 298.15 ± 0.05 K by using a jacketed glass beaker with circulating water. To minimize the concentration and temperature gradients, a magnetic stirrer was employed in the beaker. A thermometer with a reading accuracy to within ± 0.05 °C was placed in the jacket beaker. The e.m.f. of both the cation and anion ion selective electrodes were measured against a double junction reference electrode immersed in the solution in the jacketed beaker. The potentiometer readings were recorded only when the drift of the response was less than 0.1 mV in about 10 minutes. The readings for the cation and the anion electrodes were recorded simultaneously. All instruments were grounded prior to and during the experiments.

2.3 Experimental Results and Discussions

The experimental measurements were carried out at 298.15 ± 0.05 K for four systems in an electrochemical cell such as

Chapter 2. ELECTROCHEMICAL MEASUREMENT OF ACTIVITY COEFFICIENTS OF AMINO ACIDS...

Potassium ISE | electrolyte (m_S) + solute (m_A) + water | reference electrode

Nitrate ISE | electrolyte (m_S) + solute (m_A) + water | reference electrode.

The activity coefficients of the electrolyte measured at various concentrations of amino acid are listed in Table 2.1 to Table 2.4. Table 2.5 lists the pooled standard deviations (Skoog and Leary, 1992), which fall within the range from 0.002 to 0.005. At least three replicates were measured for each point. The results for $\Delta E^{(1)}$ and $\Delta E^{(2)}$ are listed in Tables A-1 to A-4 in Appendix A.

For the calculation of the activity coefficients of the amino acids in the electrolyte solutions, the ratio of the activity coefficients of the electrolytes in the solutions with and without amino acids were correlated with the following equation (Khoshkbarchi and Vera, 1996a):

$$\nu \ln \frac{\gamma_{\pm}^{(2)}}{\gamma_{\pm}^{(1)}} = C_1 m_A + C_2 m_S m_A + C_3 m_A^2 + C_4 m_A m_S^2 + C_5 m_A^3 + C_6 m_S m_A^2 \quad (2-10)$$

The values of the constants and the root mean square deviation obtained in the correlation of results are given in Table 2.6. The root mean square deviation is defined as:

$$r.m.s.d. = \sqrt{\frac{\sum \left[\left(\gamma_{\pm}^{(2)} \gamma_{\pm}^{(1)} \right)^{cal} - \left(\gamma_{\pm}^{(2)} \gamma_{\pm}^{(1)} \right)^{exp} \right]^2}{n}} \quad (2-11)$$

As shown in Table 2.6, the six parameters equation 2.10 gives a very accurate correlation necessary for further calculations.

According to thermodynamics, the activity coefficient of amino acid is related to the mean ionic activity coefficient of the electrolyte via the cross differential relation (Lewis and Randall, 1961):

Chapter 2. ELECTROCHEMICAL MEASUREMENT OF ACTIVITY COEFFICIENTS OF AMINO ACIDS...

Table 2.1 Experimental results for NaNO₃ + DL-serine system

m NaNO ₃ ⇒	0.1	0.3	0.5	0.7	1.0
m DL-serine ⇓	$\gamma_{\pm}^{(2)} / \gamma_{\pm}^{(1)}$				
0	1	1	1	1	1
0.05	0.9889	0.9915	0.9935	0.9948	0.9961
0.1	0.9793	0.9844	0.9870	0.9889	0.9922
0.15	0.9703	0.9767	0.9806	0.9851	0.9870
0.2	0.9621	0.9697	0.9754	0.9793	0.9831
0.25	0.9540	0.9634	0.9703	0.9742	0.9774
0.3	0.9459	0.9558	0.9628	0.9691	0.9729
0.35	0.9385	0.9496	0.9577	0.9646	0.9691
0.4	0.9312	0.9428	0.9527	0.9584	0.9659

Table 2.2 Experimental results for NaNO₃ + DL-valine system

m KNO ₃ ⇒	0.1	0.3	0.5	0.7	1.0
m DL-valine ⇓	$\gamma_{\pm}^{(2)} / \gamma_{\pm}^{(1)}$				
0	1	1	1	1	1
0.1	0.9973	0.9980	1.0006	1.0027	1.0040
0.2	0.9921	0.9960	0.9999	1.0053	1.0079
0.3	0.9868	0.9934	0.9993	1.0033	1.0106
0.4	0.9816	0.9889	0.9974	1.0040	1.0126
0.5	0.9751	0.9869	0.9987	1.0040	1.0126

Chapter 2. ELECTROCHEMICAL MEASUREMENT OF ACTIVITY COEFFICIENTS OF AMINO ACIDS...

Table 2.3 Experimental results for KNO₃ + DL-serine system

m KNO ₃ ⇒	0.1	0.3	0.5	0.7
m DL-serine ↓	$\gamma_{\pm}^{(2)} / \gamma_{\pm}^{(1)}$			
0	1	1	1	1
0.01	0.9962	0.9977	0.9992	1
0.03	0.9940	0.9955	0.9977	1
0.05	0.9910	0.9932	0.9962	0.9970
0.1	0.9814	0.9858	0.9925	0.9940
0.15	0.9703	0.9806	0.9873	0.9917
0.2	0.9594	0.9747	0.9828	0.9888
0.25	0.9501	0.9681	0.9784	0.9851
0.3	0.9415	0.9623	0.9747	0.9836
0.35	0.9330	0.9558	0.9710	0.9806
0.4	0.9240	0.9493	0.9681	0.9762

Table 2.4 Experimental results for KNO₃ + glycine system

m KNO ₃ ⇒	0.1	0.3	0.5	0.7
m Glycine ↓	$\gamma_{\pm}^{(2)} / \gamma_{\pm}^{(1)}$			
0	1	1	1	1
0.1	0.9797	0.9842	0.9887	0.9917
0.3	0.9433	0.9576	0.9657	0.9745
0.7	0.8795	0.9101	0.9325	0.9439
1	0.8384	0.8795	0.9067	0.9219
1.3	0.8010	0.8513	0.8842	0.9060
1.6	0.7706	0.8270	0.8636	0.8917
2	0.7346	0.7974	0.8390	0.8682
2.3	0.7098	0.7813	0.8245	0.8545
2.6	0.6897	0.7643	0.8121	0.8422

Chapter 2. ELECTROCHEMICAL MEASUREMENT OF ACTIVITY COEFFICIENTS OF AMINO ACIDS...

Table 2.5 Pooled standard deviation of the measured ratio $\gamma_{\pm}^{(2)} / \gamma_{\pm}^{(1)}$

System	Number of data points	pooled standard deviation
A) NaNO ₃ + DL-serine + H ₂ O	120	0.002
B) NaNO ₃ + DL-valine + H ₂ O	75	0.002
C) KNO ₃ + DL-serine + H ₂ O	120	0.003
D) KNO ₃ + Glycine + H ₂ O	108	0.005

Table 2.6 Coefficients of eqs.(2-10) and (2-13)

System ⁺	C ₁	C ₂	C ₃	C ₄	C ₅	C ₆	100× r.m.s.d
A	-0.461178	0.475671	0.186180	-0.183626	-0.054472	-0.194646	0.0058
B	-0.084055	0.278430	-0.159087	-0.083077	-0.162574	-0.038425	0.0776
C	-0.490994	0.941036	-0.029429	-0.550696	0.142561	-0.088478	0.0719
D	-0.457793	0.622482	0.053941	-0.320274	-0.002268	-0.043574	0.1518

r.m.s.d = root mean square deviation defined by eq.(2-11).

+ systems A, B, C, D are those listed in Table 2.5

Chapter 2. ELECTROCHEMICAL MEASUREMENT OF ACTIVITY COEFFICIENTS OF AMINO ACIDS...

$$\left(\frac{\partial \ln \gamma_A}{\partial m_S} \right)_{m_A, T, P} = \nu \left(\frac{\partial \ln \gamma_{\pm}}{\partial m_A} \right)_{m_S, T, P} \quad (2-12)$$

Combining eqs.(2.10) and (2.12), the activity coefficient of amino acid is given by:

$$\ln \frac{\gamma_A^{(2)}}{\gamma_A^{(1)}} = C_1 m_S + \frac{1}{2} C_2 m_S^2 + 2C_3 m_A m_S + \frac{1}{3} C_4 m_S^3 + 3C_5 m_A^2 m_S + C_6 m_S^2 m_A \quad (2-13)$$

In fact, eq.(2.10) is a virial expansion and the first term, C_1 , represents the pairwise interaction between amino acid molecules and electrolyte molecules. For all the 4 systems studied here, this term is negative, which means an attractive interaction (Khoshkbarchi and Vera, 1996a). The presence of electrolytes decreases the activity of amino acids. Figure 2.2 to 2.5 illustrate the correlation and calculation results obtained using eqs.(2.10) and (2.13).

Figure 2.2 depicts the effect of DL-serine and NaNO_3 concentrations on the ratios of activity coefficients $\gamma_{\pm}^{(2)}/\gamma_{\pm}^{(1)}$ and $\gamma_A^{(2)}/\gamma_A^{(1)}$. As shown in the figure, both $\gamma_{\pm}^{(2)}/\gamma_{\pm}^{(1)}$ and $\gamma_A^{(2)}/\gamma_A^{(1)}$ decrease as the molalities of NaNO_3 and DL-serine increase. The presence of the amino acid decreases the mean ionic activity coefficient.

Figure 2.3 depicts the effect of DL-serine and KNO_3 concentration on $\gamma_{\pm}^{(2)}/\gamma_{\pm}^{(1)}$ and $\gamma_A^{(2)}/\gamma_A^{(1)}$. The behavior of the activity coefficients is similar to the systems illustrated in Fig.2.2. Both $\gamma_{\pm}^{(2)}/\gamma_{\pm}^{(1)}$ and $\gamma_A^{(2)}/\gamma_A^{(1)}$ decrease as the molalities of NaNO_3 and DL-serine increase. At high concentration of electrolyte, $\gamma_{\pm}^{(2)}/\gamma_{\pm}^{(1)}$ decreases less with the increase in concentration of DL-serine for the case of KNO_3 than for the case of NaNO_3 . In the figures this effect is somewhat obscured by the need to use different scales.

Chapter 2. ELECTROCHEMICAL MEASUREMENT OF ACTIVITY COEFFICIENTS OF AMINO ACIDS...

Figure 2.2 Effect of concentrations of NaNO₃ and DL-serine, in aqueous solutions at 298.2 K, on the ratios $\gamma_z^{(2)}/\gamma_z^{(1)}$; $\gamma_A^{(2)}/\gamma_A^{(1)}$. Dots: experimental points; curves: equations (2-10), (2-13), respectively.

Chapter 2. ELECTROCHEMICAL MEASUREMENT OF ACTIVITY COEFFICIENTS OF AMINO ACIDS...

Figure 2.3 Effect of concentrations of KNO_3 and DL-serine, in aqueous solutions at 298.2 K, on the ratios $\gamma_z^{(2)}/\gamma_z^{(1)}$; $\gamma_A^{(2)}/\gamma_A^{(1)}$. Dots: experimental points; curves: equations (2-10), (2-13), respectively.

Chapter 2. ELECTROCHEMICAL MEASUREMENT OF ACTIVITY COEFFICIENTS OF AMINO ACIDS...

In both cases, the effect of the amino acid decreases as concentration of the electrolyte increases. The similarity of these two systems is natural since the only difference between them is the anion K^+/Na^+ . This effect suggests that higher electrolyte concentrations screen more the electrostatic ion-dipole interactions between DL-serine and the ions, and also increase the importance of the short range interactions, resulting in a smaller effect of a change of DL-serine concentration on the mean ionic activity coefficient. Similarly there is a smaller effect of changes in electrolyte concentration on the activity coefficient of the amino acid.

Figure 2.4 depicts the effect of DL-valine and $NaNO_3$ concentrations on the ratio of activity coefficients, $\gamma_{\pm}^{(2)}/\gamma_{\pm}^{(1)}$ and $\gamma_A^{(2)}/\gamma_A^{(1)}$. At high concentration of $NaNO_3$ (1.0 m), $\gamma_{\pm}^{(2)}/\gamma_{\pm}^{(1)}$ increases as the molality of DL-valine increases, while at concentration of $NaNO_3$ around 0.7, $\gamma_{\pm}^{(2)}/\gamma_{\pm}^{(1)}$ decreases as the molality of DL-valine increases. For the activity coefficients of the amino acid (DL-valine), a minimum value (against the concentration of $NaNO_3$) occurs at each particular concentration of amino acid. When the concentrations of DL-valine are below 0.5, $\gamma_A^{(2)}/\gamma_A^{(1)}$ decreases as the concentration of electrolyte increases, while at 0.5 m the values of $\gamma_A^{(2)}/\gamma_A^{(1)}$ become less than those at 0.3 m. A similar behavior was observed for the DL-alanine + KCl + H_2O system (Soto-Campos et al. 1997).

In fact the overlap of lines is due to the presentation of three dimensional data in a two-dimensional plot. Additional information is gained by re-plotting the values of $\gamma_A^{(2)}/\gamma_A^{(1)}$ as a function of m_A for fixed values of m_S . Such a plot is presented in Figure 2.5. In this case, the variation of the trace activity coefficient of the amino acid in salt solutions is explicitly shown as a function of salt molality.

The difference between the effect in DL-serine + $NaNO_3$ system and that in DL-valine + $NaNO_3$ system may be a result of the difference of the molecular structures of these two amino acids. In DL-serine there is a -OH group in its hydrocarbon backbone. The presence of this -OH group decreases the hydrophobic interactions between the

Chapter 2. ELECTROCHEMICAL MEASUREMENT OF ACTIVITY COEFFICIENTS OF AMINO ACIDS...

Figure 2.4 Effect of concentrations of NaNO₃ and DL-valine, in aqueous solutions at 298.2 K, on the ratios $\gamma_z^{(2)}/\gamma_z^{(1)}$; $\gamma_A^{(2)}/\gamma_A^{(1)}$. Dots: experimental points; curves: equations (2-10), (2-13), respectively.

Chapter 2. ELECTROCHEMICAL MEASUREMENT OF ACTIVITY COEFFICIENTS OF AMINO ACIDS...

hydrocarbon backbone of DL-serine and the molecules of water. Besides, at high concentrations of DL-valine and NaNO_3 , probably as a result of the formation of the ion-pair complexes, the electrostatic interactions decrease, leading to an increase in the activity coefficient of DL-valine with an increase of NaNO_3 molality.

Figure 2.6 depicts the effect of glycine and KNO_3 concentrations on $\gamma_{\pm}^{(2)}/\gamma_{\pm}^{(1)}$ and $\gamma_A^{(2)}/\gamma_A^{(1)}$. The behavior of activity coefficients for this system is similar to DL-serine + NaNO_3 + H_2O as discussed above. But the activity coefficient decreases more in the former system than in the latter. This difference may be due to the difference of the structure of DL-serine and glycine: DL-serine has a $-\text{OH}$ group in its hydrocarbon backbone and has one $-\text{CH}_2$ group more than glycine. The formation of ion-pair complexes in the glycine system and the effect of $-\text{OH}$ may both contribute to the difference.

Figure 2.5 Re-plot of $\gamma_A^{(2)}/\gamma_A^{(1)}$ as a function of m_A for fixed values of m_S

Chapter 2. ELECTROCHEMICAL MEASUREMENT OF ACTIVITY COEFFICIENTS OF AMINO ACIDS...

Figure 2.6 Effect of concentrations of KNO₃ and glycine, in aqueous solutions at 298.2 K, on the ratios $\gamma_{\pm}^{(2)}/\gamma_{\pm}^{(1)}$; $\gamma_A^{(2)}/\gamma_A^{(1)}$. Dots: experimental points; curves: equations (2-10), (2-13), respectively.

CHAPTER 3: MEASUREMENT OF ACTIVITY COEFFICIENTS USING THE ISOPIESTIC METHOD

3.1 Motivation

For the activity coefficients in $\text{KNO}_3 + \text{DL-serine} + \text{H}_2\text{O}$ system, the results obtained in this work are different from previous measurements obtained by a Summer Student using the EC method (Silverberg, 1998). The activity coefficients of DL-serine obtained by Silverberg were greater than 1.0 when KNO_3 molality was greater than 0.3.

In the calibration of the slope (S) for eq.(2-9) using eq.(2-7) and data of the system $\text{KNO}_3 + \text{water}$, it was found that the response of the pair of electrodes was slightly nonlinear. This is possibly due to interferences when both potassium ISE and nitrate ISE are used in the measurement. The details of this phenomenon and a theoretical explanation can be found in literature (Haghtalab, 1990 and references cited in that work). Some error could be introduced by this interference in the measurement recorded by both electrodes.

With the above motivation, the purpose of this part of work was to use the isopiestic method to measure the activity coefficients for this system and compare the results with those obtained from the EC method.

3.2 Theoretical Background

When a ternary system, amino acid (m_A) + electrolyte (m_S) + water, is in equilibrium with a reference system (electrolyte + water), then the activities of water in the two systems are equal:

$$a_W = a_W^{\text{Ref}} \quad (3-1)$$

According to the definition of osmotic coefficient ϕ_s for the electrolyte solution:

Chapter 3. MEASUREMENT OF ACTIVITY COEFFICIENT USING THE ISOPIESTIC METHOD

$$55.51 \ln a_w = -\nu m_s \phi_s \quad (3-2)$$

eq.(3-1) can be written as:

$$m_s \phi_s = m_s^{\text{Ref}} \phi_s^{\text{Ref}} \quad (3-3)$$

If the osmotic coefficient of the reference solution is known as a function of molality of the electrolyte (m_s), then eq.(3-3) can be used to calculate the unknown ϕ_s . For a solution containing one solute (say electrolyte), the mean ionic activity coefficient can be calculated with the following equation:

$$\ln \gamma_{\pm} = (\phi_s - 1) + \int_0^{m_s} (\phi_s - 1) d \ln m_s \quad (3-4)$$

For a solute A in a ternary system, the cross differential relation should be used (Bower and Robinson, 1965):

$$\left(\frac{\partial \ln \gamma_A}{\partial m_s} \right)_{m_A} = \nu \left(\frac{\partial \ln \gamma_{\pm}}{\partial m_A} \right)_{m_s} = \frac{\Delta}{m_A m_s} \quad (3-5)$$

where Δ is defined as:

$$\Delta = \nu m_s \phi_s - \nu m_s \phi_s^0 - m_A \phi_A^0 \quad (3-6)$$

where ϕ_s^0 represents the osmotic coefficient of the binary solution (electrolyte + water) at molality m_s , which is not equal to ϕ_s^R (at molality m_s^R), and ϕ_A^0 , the osmotic coefficient of the binary solution (amino acid + water) at molality m_A .

At each set of molalities, (m_s , m_A), Δ is evaluated, and a polynomial functional dependence is assumed:

Chapter 3. MEASUREMENT OF ACTIVITY COEFFICIENT USING THE ISOPIESTIC METHOD

$$\frac{\Delta}{m_A m_S} = C_0 + C_1 m_A + C_2 m_S + C_3 m_A^2 + C_4 m_S^2 + C_5 m_A m_S \quad (3-7)$$

The coefficients of eq.(3-7) are obtained by fitting the experimental data with the equation. Then the activity coefficients are calculated with the following equations (Bower and Robinson, 1965):

$$\ln \frac{\gamma_A^{(2)}}{\gamma_A^{(1)}} = m_S \left(C_0 + C_1 m_A + \frac{1}{2} C_2 m_S + C_3 m_A^2 + \frac{1}{3} C_4 m_S^2 + \frac{2}{3} C_5 m_A m_S \right) \quad (3-8)$$

$$\nu \ln \frac{\gamma_{\pm}^{(2)}}{\gamma_{\pm}^{(1)}} = m_A \left(C_0 + \frac{1}{2} C_1 m_A + C_2 m_S + \frac{1}{3} C_3 m_A^2 + C_4 m_S^2 + \frac{2}{3} C_5 m_A m_S \right) \quad (3-9)$$

3.3 Experimental Set Up and Measurements

Several versions of set up were built and tested before the final one was accepted.

The first set up used was constructed as follows: four de-connectable flasks with volume 100 ml each were connected (ground glass connection) with a straight glass tube closed by a vacuum valve. It took too a very long time to reach the equilibrium.

The second version of the set up was different from the first one: four flasks, of 100 ml volume, were connected to a spherical chamber to avoid the diffusion through the tube. Still it was too difficult to reach equilibrium and it was decided to shorten the connecting necks of the flasks and decrease their volume.

In the third version of the set up four cells with volume of around 10 ml each were connected permanently with a spherical chamber. This set up worked much better than previous versions. Unfortunately it was difficult to fill the solutions through the upper opening of the chamber.

Finally the fourth and last version was constructed: four cells with 10 ml volume were connected with a semi-spherical chamber using screw connections

Chapter 3. MEASUREMENT OF ACTIVITY COEFFICIENT USING THE ISOPIESTIC METHOD

(CG-350, CHEMGLASS), as illustrated in Figure 3.1. While the three previous setups were thermostabilized in a water bath, the fourth setup was put in a custom-made air bath which kept the temperature at 298.15 ± 0.05 K. The solutions were distributed in the four cells as follows: two cells contained binary solution ($\text{KNO}_3 + \text{H}_2\text{O}$), and two cells contained the ternary solution ($\text{DL-serine} + \text{KNO}_3 + \text{H}_2\text{O}$). The concentrations of the solutions were measured using the Atomic Absorption (AA) (Thermo Jarrell Ash, Smith-Hieftje 11) method. Generally, around 4 weeks were needed for the system to reach equilibrium.

Operation procedure

- (1) Two reference solutions ($\text{KNO}_3 + \text{water}$) with the different concentrations of the electrolyte were filled in two cells. The other two cells were filled with solutions of the unknown system ($\text{KNO}_3 + \text{DL-serine} + \text{water}$) with the same salt and amino acid concentration ratio, but different amount of water.
- (2) The chamber was evacuated and the air bath was set at 298.15 K.
- (3) The concentrations in each cell were measured periodically. When the concentrations of the two reference systems were found to be equal and the concentrations of the two unknown systems were also equal, the system was considered to be in equilibrium.

Chapter 3. MEASUREMENT OF ACTIVITY COEFFICIENT USING THE ISOPIESTIC METHOD

1: vacuum stopcock; 2: vapor chamber; 3: sealing washer; 4: cell (total 4×10 ml); 5: magnet; 6: chem-thread (CG-350, CHEMGLASS)

Figure 3.1 View of isopiestic set up

Chapter 3. MEASUREMENT OF ACTIVITY COEFFICIENT USING THE ISOPIESTIC METHOD

3.4 Results and Discussions

The values for osmotic coefficients in the binary system (KNO_3 + water; DL-serine + water) at various solute concentrations were obtained as follows. First the activity coefficients of amino acid and electrolyte in aqueous solutions at 298.15 K were obtained from literature (Fasman, 1976; Zaytsev and Aseyev, 1992), respectively. For the mean ionic activity coefficient of the electrolyte, the extended Debye-Huckel equation for 1:1 electrolyte is given by (Butler, 1998):

$$\ln \gamma_{\pm} = -\frac{A\sqrt{m}}{1 + B_1\sqrt{m}} + B_2 m + B_3 m^2 + \dots \quad (3-10)$$

Theoretical value A is 1.174 at 298.15 K (Butler, 1998). In the Davies model (Davies, 1962), only the first two terms are used with $B_1 = 1$ and $B_2 = -0.2$. In this case the correlation error was considered high with AAD (average absolute deviation) 1.23%. For a better correlation, three terms were used, and treated both B_1 and B_3 were treated as adjustable parameters. The values of the parameters and the correlation results are listed in Table 3.1.

For the asymmetrically normalized activity coefficient of the amino acid, we used the following polynomial function:

$$\ln \gamma_A = b m_A + c m_A^2 + d m_A^3 \quad (3-11)$$

The values of the parameters and the correlation results are listed in Table 3.2.

Table 3.1 Correlation results by eq.(3-10)

System	B_1	B_2	B_3	AAD%
KNO_3 + water	0.8927	-0.2	0.009807	0.55

Chapter 3. MEASUREMENT OF ACTIVITY COEFFICIENT USING THE ISOPIESTIC METHOD

Table 3.2 Correlation results by eq.(3-11)

System	<i>b</i>	<i>c</i>	<i>d</i>	AAD%
DL-serine + water	-0.263674	0.052884	-0.004677	0.12

The osmotic coefficient is given by (Robinson and Stokes, 1959):

$$\phi = 1 + \frac{1}{m_0} \int_0^m m \, d \ln \gamma \quad (3-12)$$

For electrolyte in water, from eqs.(3-10) and (3-12), we have

$$\phi = 1 - \frac{A\sqrt{m}}{(B_1\sqrt{m})^3} \left[1 + B_1\sqrt{m} - 2 \ln(1 + B_1\sqrt{m}) - \frac{1}{1 + B_1\sqrt{m}} \right] + \frac{1}{2} B_2 m + \frac{2}{3} B_3 m^2 \quad (3-13)$$

and for DL-serine in water, from eqs.(3-11) and (3-12):

$$\phi = 1 + \frac{1}{2} b m_{,1} + \frac{2}{3} c m_{,1}^2 + \frac{3}{4} d m_{,1}^3 \quad (3-14)$$

Eqs.(3-13) and (3-14) were used to calculate the osmotic coefficients in eq.(3-6) for the two aqueous systems at various concentrations of solutes, respectively.

Finally, after obtaining the coefficients by fitting eq.(3-7), eqs.(3-8) and (3-9) can be used to calculate the activity coefficients for the ternary systems. The measured data are listed in Table 3.2, and the regressed values for the coefficient (C_0 to C_5) are given in Table 3.3.

3.5 Comparison with the Electrochemical Method

Figure 3.2 depicts a comparison of the mean ionic activity coefficients obtained by the EC method and isopiestic method at the salt molality of 0.3. Figure 3.3 depicts the activity coefficients of DL-serine in the ternary system measured by two methods at amino acid molality of 0.3. Both methods give comparable trends in the sense that the activity coefficients are less than 1.0, though some differences are observed.

The deviation of the results of the IP method from those of the EC method comes from two major sources: (1) the time was still not long enough to reach the equilibrium; (2) errors from the concentration analysis. Since for the systems considered the activity coefficients do not differ significantly from 1.0 any small deviation can not be ignored for an accurate measurement.

In addition, the concentration range for both the salt and the amino acid were not wide enough: at around 0.3 in molality of the amino acid, the extrapolation of the results to lower and higher concentrations should be taken with care. In order to obtain better results over wide concentration range, more experiments would be necessary.

Time constraints of the program forced us to stop these measurements. In any case, the main objective of this section was attained since the trend of change was confirmed.

Chapter 3. MEASUREMENT OF ACTIVITY COEFFICIENT USING THE ISOPIESTIC METHOD

Table 3.2 Measured data from isopiestic method

m_A	m_S	m^{ref}	ϕ_A^0	ϕ_S^0	ϕ^{ref}	$\Delta/m_A m_S$
0.2889	0.2777	0.4389	0.9710	0.8584	0.8303	-0.3545
0.2123	0.3323	0.4517	0.9787	0.8480	0.8283	-0.3270
0.1882	0.4312	0.5376	0.9812	0.8315	0.8158	-0.3031
0.2008	0.4213	0.5345	0.9799	0.8330	0.8163	-0.3076
0.2203	0.2923	0.4156	0.9779	0.8555	0.8340	-0.3469
0.2755	0.2846	0.4384	0.9724	0.8570	0.8304	-0.3521
0.2140	0.2170	0.3366	0.9785	0.8714	0.8472	-0.3715
0.2086	0.2156	0.3323	0.9791	0.8717	0.8480	-0.3677
0.2999	0.2245	0.3911	0.9700	0.8697	0.8379	-0.3861
0.3288	0.2010	0.3834	0.9671	0.8752	0.8392	-0.3982
0.2407	0.2614	0.3956	0.9759	0.8617	0.8372	-0.3656
0.2632	0.2655	0.4122	0.9736	0.8608	0.8345	-0.3631
0.3455	0.2345	0.4259	0.9654	0.8674	0.8323	-0.3876
0.1340	0.3870	0.4635	0.9866	0.8386	0.8265	-0.2915

Table 3.3 Correlation results

C_0	C_1	C_2	C_3	C_4	C_5	r.m.s.d
-0.34559	-1.02022	0.62344	1.24695	-0.74989	0.75688	0.00327

Chapter 3 MEASUREMENT OF ACTIVITY COEFFICIENT USING THE ISOPIESTIC METHOD

Figure 3.2 Comparison of activity coefficients of DL-serine by two methods

Figure 3.3 Comparison of mean ionic activity coefficient by two methods

Chapter 3 MEASUREMENT OF ACTIVITY COEFFICIENT USING THE ISOPIESTIC METHOD

The advantages and disadvantages of the two methods are listed below:

	Advantages	Disadvantages
Electrochemical	Accurate at low concentrations, less expensive, fast	System and concentration are limited by electrode
Isopiestic	Useful for medium and high concentrations	Slow to reach to equilibrium Inaccurate at low concentrations

CHAPTER 4: MODELING

In this chapter we discuss two models. The first, presented in section 4.1, is a direct application of the model proposed previously by Kohshibarchi and Vera (1996a). The second, presented in section 4.2, is a modification of a more complex model proposed by Khoshkbarchi and Vera (1996c).

4.1 Gibbs Free Energy Model with the NRTL Equation

For the modeling of aqueous solutions containing electrolytes and non-electrolytes, the equation for the excess Gibbs free energy is normally written as:

$$G^E = G_{LR}^E + G_{SR}^E \quad (4-1)$$

where the superscripts 'LR', 'SR' refer to long and short range interactions, respectively.

A model was proposed by Kohshibarchi and Vera (1996a) in which either the Wilson or the NRTL equations were used to represent the short range contributions. In this model, the activity coefficient of an electrolyte in an electrolyte + amino acid + water system is given by:

$$\ln \gamma_i^{(m)} = \ln \gamma_i^{(m),LR} + \frac{1}{\nu} \left\{ \ln \gamma_S^{(x),SR} - \lim_{x_i \rightarrow 0} (\ln \gamma_S^{(x),SR}) + \ln \left(\frac{1 + 0.001 M_w m_S}{1 + 0.001 M_w (m_S + m_A)} \right) \right\} \quad (4-2)$$

where the subscripts 'S', 'A' and 'W' refer to electrolyte, amino acid, and water; the superscripts (m) and (x) refer to molality and molar fraction based activity coefficient, respectively; M_w is the molecular weight of water, m_S and m_A refer to the molalities of electrolyte and amino acid in the solution. Clearly, at zero amino acid concentration, the mean ionic activity coefficient of the electrolyte reduces to the long range (LR) contribution. The last term of eq.(4-2) arises from the conversion of the molality based activity coefficient into the molar fraction based one. In the case of the binary system electrolyte + water, the following equation holds:

$$\gamma_{\pm}^{(x)} = \gamma_{\pm}^{(m)} (1 + 0.001 M_w m_s) \quad (4-3)$$

When values of the mean ionic activity coefficient are needed, a model for the long range interaction term, namely activity coefficient of electrolyte in water, is required. There are many different models for this purpose, such as Bromley model (Bromley, 1973), K-V model (Kohshkbarchi and Vera, 1996b). In this work, we only need the ratio of the activity coefficients of the electrolyte in a aqueous solution in the presence and absence of the amino acid, then eq.(4-2) is written as:

$$\ln \frac{\gamma_{\pm}^{(m)}}{\gamma_{\pm}^{(m),LR}} = \ln \frac{\gamma_{\pm}^{(2)}}{\gamma_{\pm}^{(1)}} = \frac{1}{\nu} \left\{ \ln \gamma_s^{(x),SR} - \lim_{x_s \rightarrow 0} (\ln \gamma_s^{(x),SR}) + \ln \left(\frac{1 + 0.001 M_w m_s}{1 + 0.001 M_w (m_s + m_A)} \right) \right\} \quad (4-4)$$

where $\gamma_{\pm}^{(m),LR}$ is the mean ionic activity coefficient of the electrolyte in the binary aqueous solution at the same electrolyte molality as in the ternary systems, which is identical to $\gamma_{\pm}^{(1)}$. Similarly, the ratios of the activity coefficients of amino acid in a aqueous solution with and without electrolyte is given by:

$$\ln \frac{\gamma_A^{(m)}}{\gamma_A^{(m),B}} = \ln \frac{\gamma_A^{(2)}}{\gamma_A^{(1)}} = \ln \gamma_A^{(x),SR} - \lim_{x_s \rightarrow 0} (\ln \gamma_A^{(x),SR}) - \lim_{x_s \rightarrow 0} (\ln \gamma_A^{(x),SR}) + \ln \left(\frac{1 + 0.001 M_w m_A}{1 + 0.001 M_w (m_s + m_A)} \right) \quad (4-5)$$

where superscript B refers to binary system. The third term in the right hand side of eq.(4-5) is introduced to meet the limit at infinite dilution (Kohshkbarchi and Vera 1996a).

As it can be seen from eq.(4-4) and (4-5), for the purposes of this work, a model for the short range interaction activity coefficient suffices to give a complete formulation. Here the NRTL equation is used:

$$\ln \gamma_i = \frac{\sum_{j=1}^3 x_j \tau_{ji} G_{ji}}{\sum_{j=1}^3 x_j G_{ji}} + \sum_{j=1}^3 \frac{x_j G_{ij}}{\sum_{k=1}^3 x_k G_{kj}} \left(\tau_{ij} - \frac{\sum_{k=1}^3 x_k \tau_{kj} G_{kj}}{\sum_{k=1}^3 x_k G_{kj}} \right) \quad (4-6)$$

with

$$G_{ij} = \exp(-\alpha \tau_{ij}) \quad (i,j = 1,2,3) \quad (4-7)$$

where τ_{ij} 's are the binary parameters, with $\tau_{ii} = 0$. Here we use the convention: 1 = S (electrolyte), 2 = A (amino acid), 3 = W (water). The binary parameters for the amino acid + water system ($\tau_{A,W}$, $\tau_{W,A}$) were estimated using literature data (Fasman, 1976). $\tau_{S,A}$ and $\tau_{S,W}$ were estimated from ternary experimental data with the assumptions $\tau_{A,S} = \tau_{S,A}$ and $\tau_{W,S} = \tau_{S,W}$. The parameter α was set constant at 0.3.

Following Kohshkbarchi and Vera (1996a) for the calculation of the short range contribution, the electrolyte was considered to be a molecular component and its molar fraction in a ternary system is calculated by:

$$x_S = \frac{n_S}{n_A + n_S + n_W} \quad (4-8)$$

4.2 A New Model Based on a Simplified Perturbation Theory

4.2.1 Theoretical Background

Khoshkbarchi and Vera (1996c) proposed a theoretical model based on the perturbation theory. For the chemical potential, using the primitive approximation, this theory gives

$$\mu_i = \mu_i^{Ref} + \mu_i^{Per} \quad (4-9)$$

where superscripts 'Ref' and 'Per' refer to the reference and perturbation contributions to the chemical potential, respectively. Using the hard sphere expression as reference model and model potentials for perturbation, the activity coefficient is expressed as

$$\ln \gamma_i = \frac{8\eta + 9\eta^2 + 3\eta^3}{(1-\eta)^3} - \frac{4\pi\rho}{kT} \left\{ \frac{8}{9} \varepsilon \sigma^3 + \frac{D^4}{9(4\pi\varepsilon_0\varepsilon_r)^2 kT \sigma^3} \right\} \quad (4-10)$$

where σ and ε are the molecular diameter and the depth of the molecular potential well according to the Lennard-Jones (LJ) potential model, respectively, ε_0 the permittivity of vacuum, ε_r , the relative dielectric constant of the medium (water), D is the dipole moment of the amino acid, $\rho = N_i/V$, the number density of the component (i) in the solution with total volume V , k is the Boltzmann constant, T is the absolute temperature. and finally, η is given by:

$$\eta = \frac{1}{6} \pi \rho \sigma^3 \quad (4-11)$$

The first term in the right hand side of eq.(4-10) represents the contribution of the reference system, which is taken as a hard sphere system, the second term is the perturbation contribution obtained by considering the dispersion forces (LJ) and dipole-dipole interaction contributions via the expression:

$$\mu_i^{Per} = 4\pi\rho \int_0^\infty u(r) g^{hs}(r) r^2 dr \quad (4-12)$$

where $u(r)$ is the interaction energy as a function of inter-molecular distances, $g^{hs}(r)$, the radial distribution function of the hard sphere reference system. A rigorous expression for $g^{hs}(r)$ can be employed to obtain an analytical equation for the perturbation contribution (Khoshkbarchi and Vera, 1998). For practical applications, however, this treatment did not improve the results. In this work, a simple step function radial distribution function was used for $g^{hs}(r)$:

$$g^{hs}(r) = \begin{cases} 0, & r < \sigma \\ 1, & r \geq \sigma \end{cases} \quad (4-13)$$

The advantages of eq.(4-10) are obvious: it is simple but has a strong theoretical background, and the modeling of results for both activity coefficients and solubility data are satisfactory. Unfortunately, dipole moments for amino acids are scarce and a quantum mechanical approach is needed to calculate this quantity, which is not convenient for engineering applications.

Khoshkbarchi and Vera (1996c) have extended the model to ternary systems. Their expression is lengthy and it is omitted here. In this work, for the ternary system we use a simpler version of the model.

4.2.2 A New Model

Binary water - amino acid system

The key quantity to derive this model is the perturbation potential. In a solution, in addition to the dispersion and the dipole-dipole interactions mentioned above, there are molecular interactions such as dipole induced interaction characterized by the polarizability, $\bar{\alpha}$, etc. A potential function, which considers all the three contributions in the primitive approximation, can be written as (Maitland et al., 1981):

$$u(r) = 4\epsilon \left[\left(\frac{\sigma}{r} \right)^{12} - \left(\frac{\sigma}{r} \right)^6 \right] - \frac{2}{(4\pi\epsilon_0\epsilon_r)^2} \left(\frac{D^4}{3kTr^6} + \frac{D^2\bar{\alpha}}{r^6} + \dots \right) \quad (4-14)$$

where the first term is the Lennard-Jones potential for the dispersion interaction.

From eqs.(4-12) to (4-14), an expression for μ_i^{Per} can be obtained, in which three terms are temperature-independent. For simplicity, the temperature-dependent terms can be neglected and combining the temperature-independent terms as shown in Appendix B, the perturbation contribution can be simply expressed as:

$$\ln \gamma_i^{Per} = -B \frac{4\pi}{kT} (\rho \sigma^3 \varepsilon) \quad (4-15)$$

where constant B includes D , $\bar{\alpha}$ and other constants. It was found that for an amino acid in aqueous solutions, the constant B can be assigned the universal value $B = 0.2$.

Therefore, for a binary amino acid + water system, the simplified model used in this work gives the following expression for the activity coefficient:

$$\ln \gamma_i = \frac{8\eta + 9\eta^2 + 3\eta^3}{(1-\eta)^3} - \frac{0.8\pi}{kT} \rho \sigma^3 \varepsilon \quad (4-16)$$

In this equation, molecules are characterized only by two parameters: ε and σ and no additional property is needed.

Ternary amino acid - electrolyte - water system

The extension of this model to ternary systems is straightforward (Kohshkbarchi and Vera 1996c). For the amino acid, the activity coefficient in the system is written as:

$$\ln \gamma_A = \ln \gamma_A^{HS} + \ln \gamma_A^{Per} \quad (4-17)$$

and for electrolyte:

$$\ln \gamma_{\pm} = \ln \gamma_{\pm}^{LR} + \frac{1}{\nu} \left[\ln \gamma_S^{HS} + \ln \gamma_S^{Per} - \lim_{\rho_i \rightarrow 0} (\ln \gamma_S^{HS} + \ln \gamma_S^{Per}) \right] \quad (4-18)$$

where the contribution to the mean ionic activity coefficient arising from the long range is exactly the value of the mean ionic activity coefficient in the binary electrolyte + water system at the same electrolyte molality as in the ternary system. Thus, $\ln \gamma_{\pm}^{LR}$ is equal to $\ln \gamma_{\pm}^{(1)}$, and with the notation used here, eq.(4-18) can be rewritten as:

$$\ln \frac{\gamma_{\pm}^{(2)}}{\gamma_{\pm}^{(1)}} = \frac{1}{\nu} \left[\ln \gamma_S^{HS} + \ln \gamma_S^{Per} - \lim_{\rho_s \rightarrow 0} (\ln \gamma_S^{HS} + \ln \gamma_S^{Per}) + \ln \left(\frac{1 + 0.001 M_w m_s}{1 + 0.001 M_w (m_s + m_A)} \right) \right] \quad (4-19)$$

where, as discussed previously, the last term is introduced to convert the activity coefficient from a molar fraction scale into a molality scale. Similarly, for the amino acid, eq.(4-17) is written as:

$$\ln \frac{\gamma_A^{(2)}}{\gamma_A^{(1)}} = \ln \gamma_A^{HS} + \ln \gamma_A^{Per} - \lim_{\rho_s \rightarrow 0} (\ln \gamma_A^{HS} + \ln \gamma_A^{Per}) + \ln \left(\frac{1 + 0.001 M_w m_A}{1 + 0.001 M_w (m_s + m_A)} \right) \quad (4-20)$$

Various expressions for the activity coefficients of mixture of hard spheres are available in literature. For instance, Taghikhani et al. (1999) recently proposed an expression which leads to improved results for the radial distribution function. For simplicity, in this work the expression proposed by Mansoori et al. (1971) was employed:

$$\ln \gamma_i^{HS} = -\ln(1 - \eta_3) + D + E + 3F \quad (4-21)$$

with

$$D = \frac{\eta_0 \sigma_i^3}{1 - \eta_3} + \frac{3\eta_1 \eta_2 \sigma_i^3}{(1 - \eta_3)^2} + \frac{\eta_2^3 \sigma_i^3 (3 - \eta_3)}{(1 - \eta_3)^3} \quad (4-22)$$

$$E = \frac{3\eta_2 \sigma_i + 3\eta_1 \sigma_i^2}{1 - \eta_3} + \frac{9\eta_2^2 \sigma_i^2}{2(1 - \eta_3)^2} - \frac{\eta_2^3 \sigma_i^3}{\eta_3^3} \left[2\ln(1 - \eta_3) + \frac{\eta_3(2 - \eta_3)}{1 - \eta_3} \right] \quad (4-23)$$

$$F = \frac{\eta_2^2 \sigma_i^2}{\eta_3^2} \left[\ln(1 - \eta_3) + \frac{\eta_3}{1 - \eta_3} - \frac{\eta_3^2}{2(1 - \eta_3)^2} \right] \quad (4-24)$$

where

$$\eta_n = \frac{\pi}{6} \sum_{k=1}^3 \rho_k \sigma_k^n \quad (n = 0, 1, 2, 3) \quad (4-25)$$

Chapter 4. MODELING

For the case of an amino acid + electrolyte + water system, in the primitive approach, only the amino acid and the electrolyte contribute to the sum.

Now we need an expression for the perturbation term. For this purpose, eq.(4-15) is extended to ternary systems. The chemical potential based on the first order Barker-Henderson perturbation theory is given by (Tiepel and Gubbins, 1973):

$$\mu_i^{per} = 2\pi \frac{\partial}{\partial \rho_i} \left[\sum_{i=1}^c \sum_{j=1}^c \rho_i \rho_j \int_{\sigma_{ij}}^{\infty} u_{ij}(r) g_{ij}^{HS}(r) r^2 dr \right] \quad (4-26)$$

where c is the total number of components. The following mixing rules are used for the molecular size and energy parameters:

$$\sigma_{AS} = \sigma_{SA} = \frac{1}{2}(\sigma_A + \sigma_S)(\delta^{(0)} + \delta^{(1)} \rho_A \rho_S) \quad (4-27)$$

$$\epsilon_{AS} = \epsilon_{SA} = \sqrt{\epsilon_A \epsilon_S} \quad (4-28)$$

The two constants ($\delta^{(0)}$ and $\delta^{(1)}$) are adjustable parameters to be obtained from the measured activity coefficients data for the ternary system. As shown in appendix C, the perturbed activity coefficients for amino acid and electrolyte are given by eq.(4-29) and (4-30), respectively.

$$\ln \gamma_A^{per} = -\frac{0.8\pi}{kT} \left[\rho_A \epsilon_{AA} \sigma_{AA}^3 + \rho_S \epsilon_{AS} \sigma_{AS}^3 + 3\rho_A \rho_S \epsilon_{AS} \sigma_{AS}^2 \left(\frac{\partial \sigma_{AS}}{\partial \rho_A} \right) \right] \quad (4-29)$$

$$\ln \gamma_S^{per} = -\frac{0.8\pi}{kT} \left[\rho_S \epsilon_{SS} \sigma_{SS}^3 + \rho_A \epsilon_{AS} \sigma_{AS}^3 + 3\rho_A \rho_S \epsilon_{AS} \sigma_{AS}^2 \left(\frac{\partial \sigma_{AS}}{\partial \rho_S} \right) \right] \quad (4-30)$$

where:

$$\frac{\partial \sigma_{AS}}{\partial \rho_A} = \frac{1}{2}(\sigma_A + \sigma_S) \delta^{(1)} \rho_S \quad (4-31)$$

$$\frac{\partial \sigma_{AS}}{\partial \rho_S} = \frac{1}{2}(\sigma_A + \sigma_S) \delta^{(1)} \rho_A \quad (4-32)$$

As shown in Appendix D, the activity coefficients given by eqs.(4-29) and (4-30) satisfy the following cross differential relation:

$$\left(\frac{\partial \ln \gamma_A^{Per}}{\partial \rho_S} \right)_{T, \rho_1} = \left(\frac{\partial \ln \gamma_S^{Per}}{\partial \rho_A} \right)_{T, \rho_S} \quad (4-33)$$

The molecular parameters for electrolyte are calculated from those of the ions by the following mixing rules (Khoshkbarchi and Vera, 1997):

$$\varepsilon_S = \sqrt{\varepsilon_+ \varepsilon_-} \quad (4-34)$$

$$\sigma_S = \frac{1}{2}(\sigma_+ + \sigma_-) \quad (4-35)$$

The values of parameters of ions and electrolytes used in this work are listed in Table 4.1. For NO_3^- , the energy parameter was calculated with the following equation (Khoshkbarchi and Vera, 1997):

$$\varepsilon_i/k = 2.2789 \times 10^{-11} \bar{\alpha}_i^{1.5} \eta_i^{0.5} / \sigma_i^6 \quad (4-36)$$

where the polarizability ($\bar{\alpha}$) and total number of electrons (η_i) equal $4.37 \times 10^{-30} \text{ (m}^3\text{)}$ (Zaytev and Aseyev, 1992) and 32, respectively.

Table 4.1 Values of pure component parameters of ions and electrolytes

	Na^+	K^+	NO_3^-	KNO_3	NaNO_3
$\sigma \text{ (}\times 10^{-10} \text{ m)}$	1.90 ⁽¹⁾	2.66 ⁽¹⁾	4.28 ⁽²⁾	3.47	3.09
$\varepsilon/k \text{ (K)}$	96.0 ⁽¹⁾	214.0 ⁽¹⁾	191.6	202.5	135.6

(1): Khoshkbarchi and Vera (1997); (2) Zaytev and Aseyev, (1992).

4.3 Calculation Results

4.3.1 Binary systems

For testing the new model, eq.(4-16) was used to correlate the activity coefficient of amino acid in aqueous solution. The two parameters: σ and ε , were regressed from experimental data (Fasman, 1976). According to the definition of molality of amino acid, m_A , the number density used in the equation is calculated by the following equation:

$$\rho_A = 1000m_A N_a \quad (4-37)$$

The activity coefficient based on molality is changed into the one based on molar fraction by the following equation:

$$\gamma_A^{(x)} = \gamma_A^{(m)} (1 + 0.001M_A m_A) \quad (4-38)$$

The objective function used is given by:

$$OF. = \sum_i (\gamma_A^{\text{exp}} - \gamma_A^{\text{cal}})^2 \quad (4-39)$$

Table 4-2 lists the results for 7 systems. Figure 4.1 depicts the results for 4 systems.

Table 4.2 Calculation results for activity coefficients of amino acids

System	ε/k (K)	$\sigma \times 10^{-10}$ (m)	s.e.e
Alanine	365.9	3.94	0.0024
α -amino n-butyric acid	341.8	4.14	0.0046
Glycine	1020.0	4.25	0.0094
Proline	231.8	3.84	0.0285
DL-serine	1205.0	4.28	0.0099
Valine	321.5	5.32	0.0066
L-hydroxyproline	441.8	3.57	0.0021

The standard error of estimate is defined as (Draper and Smith, 1981):

$$s.e.e = \sqrt{\sum_i (\gamma_A^{\text{cal}} - \gamma_A^{\text{exp}})^2 / (n_D - 2)}.$$

Figure 4.1 Correlation results for activity coefficients of amino acids in binary aqueous solutions

4.3.2 Ternary systems

Both the Gibbs free energy model (Kohshibarchi and Vera, 1996a) and the new simplified perturbation model were used to correlate the 4 ternary systems measured in this work. The objective function used to regress the parameters is defined as:

$$O.F. = \sum_i \left[\left(\gamma_z^{(2)} / \gamma_z^{(1)} \right)_i^{\text{exp}} - \left(\gamma_z^{(2)} / \gamma_z^{(1)} \right)_i^{\text{cal}} \right]^2 \quad (4-40)$$

Table 4.3 lists the parameters and the results obtained using the Gibbs free energy model.

Table 4.3 Results for ternary systems by Gibbs free energy model

System	$\tau_{A,S}$	$\tau_{S,W}$	$\tau_{A,W}$	$\tau_{W,A}$	s.e.e
glycine-KNO ₃ -H ₂ O	-0.0472	0.0726	-1.9610	4.5801	0.0341
DL-serine-NaNO ₃ -H ₂ O	-0.5607	0.3311	-2.0771	3.8481	0.0067
DL-serine-KNO ₃ -H ₂ O	-0.5166	0.3042	-2.0771	3.8481	0.0106
DL-valine-NaNO ₃ -H ₂ O	-0.8756	0.5346	-0.8496	1.4371	0.0092

NRTL parameters: $\tau_{A,W}$ and $\tau_{W,A}$ from amino acid + water systems (Fasman, 1976);
 $\tau_{A,S}$ and $\tau_{S,W}$ from ternary systems (this work).

$$s.e.e = \sqrt{\frac{\sum_i \left\{ \left[\left(\gamma_z^{(2)} / \gamma_z^{(1)} \right)_i^{\text{exp}} - \left(\gamma_z^{(2)} / \gamma_z^{(1)} \right)_i^{\text{cal}} \right]^2 \right\}}{n_D - 2}}$$

Table 4.4 lists the results obtained using the simplified perturbation model. The parameters used for the amino acids are those from Table 4.2, and for the electrolytes those from Table 4.1. Figure 4.2 depicts the results for Glycine+KNO₃+water system obtained with the new simplified perturbation model using two adjustable parameters. The results obtained with keeping one or two parameters constant, and also those obtained with the Gibbs free energy model, are almost not distinguishable from those presented when plotted in the figure.

For testing the predictive features of the model, the two parameters were taken as fixed values for all systems: $\delta^{(0)} = 1.6$ and $\delta^{(1)} = -0.1 \times N_a^2 \times 10^6$. The root mean square deviations are also listed in Table 4.4. As is seen from the table, the prediction results are acceptable and comparable with the results obtained from the Gibbs free energy model.

Table 4.4 Results for ternary systems by perturbation model

System	$\delta^{(0)}$	$\delta^{(1)}/N_a^2 \times 10^6$	s.e.e	s.e.e ^a
glycine-KNO ₃ -H ₂ O	1.4984	-0.0673	0.0115	0.0334
DL-serine-NaNO ₃ -H ₂ O	1.6572	-0.3615	0.0036	0.0066
DL-serine-KNO ₃ -H ₂ O	1.5363	-0.6855	0.0053	0.0228
DL-valine-NaNO ₃ -H ₂ O	1.5680	-0.2363	0.0076	0.0106

a: prediction results with $\delta^{(0)} = 1.6$ and $\delta^{(1)} = -0.1 \times N_a^2 \times 10^6$.

Calculations adjusting one of the parameters and keeping the other constant produced results that only slightly better than those obtained fixing the values of both parameters. As seen from Tables 4.3 and 4.4, both methods give satisfactory correlation results for the four ternary systems. Therefore, a two parameter model suffices for practical applications. The perturbation model gives somewhat better results and can be used as prediction model with acceptable accuracy.

Figure 4.2 Correlation results by the simplified perturbation model for the mean activity coefficient of KNO₃ in ternary system (KNO₃ + glycine + H₂O).

4.4 Modeling the Solubility of Amino Acid in Water

The solubility of amino acids in water is an important information for separation purposes. For an accurate correlation of this property, a reliable activity model is required. As a further application of the model proposed for binary amino acid + water system, eq.(4-16). The solubility of amino acid in water was correlated for some systems where both activity coefficient and solubility data are available (Fasman, 1976).

Theoretical background

The above modeling for the activity coefficient was performed at a constant temperature (298.15 K). But for the solubility correlation, wide temperature ranges (273.15 to 373.15 K) have to be considered. For a better representation of the temperature dependence of the activity coefficients, the effective hard sphere diameter (EHSD) approach was employed. In this approach, the molecular diameter is considered to be a function of temperature and density. For simplicity, in this work we take only the temperature effect into account. Various expressions for the EHSD are available in literature (Ben-Amotz and Herschbach, 1990). Here the following expression obtained by Ben-Amotz and Herschbach (1990) based on the molecular simulation results was adopted:

$$\sigma_e = 1.1532\sigma \left[1 + \left(\frac{T^*}{0.527} \right)^{1/2} \right]^{-1/6} \quad (4-41)$$

where the subscript 'e' refers effective hard sphere diameter. The reduced temperature is defined as:

$$T^* = \frac{T}{\epsilon/k} \quad (4-42)$$

Now the reference term of the activity coefficient is expressed as:

$$\ln \gamma_i^{\text{Ref}} = \frac{8\eta_e + 9\eta_e^2 + 3\eta_e^3}{(1 - \eta_e)^3} \quad (4-43)$$

where

$$\eta_e = \frac{1}{6} \pi \rho \sigma_e^3 \quad (4-44)$$

The perturbation term is the same as eq.(4-15). The molecular size and energy parameters are taken from Table 4.2.

For the system considered here, when it is in equilibrium with a solid phase of pure amino acid:

$$f_A^L = f_A^S \quad (4-45)$$

If the standard state of the liquid fugacity is f_A^{0L} , then eq.(4-45) can be written as:

$$x_A \gamma_A f_A^{0L} = f_A^S \quad (4-46)$$

From the definition of fugacity (Koshkbarchi and Vera, 1996d):

$$\frac{f_A^S}{f_A^{0L}} = \exp\left(\frac{\Delta s}{R} - \frac{\Delta h}{RT}\right) \quad (4-47)$$

where Δs and Δh are the change in the molar entropy and enthalpy of the amino acid, respectively, from the standard liquid state to the solid state, and R is the gas constant. Finally, the solubility of amino acid in water is estimated from the following equation:

$$x_i \gamma_i = \exp\left(\frac{\Delta s}{R} - \frac{\Delta h}{RT}\right) \quad (4-48)$$

The two quantities $\Delta s/R$ and $\Delta h/R$ are regressed from the solubility data (Fasman, 1976).

Calculation results

Eqs.(4-48) and (4-43) with (4-15) ($B = -0.2$) were used to correlate the solubility data. In eq.(4-48) at each solubility S (g/kg-solvent), the number density is calculated with:

$$\rho_A = 1000N_a S / M_A \quad (4-49)$$

where M_A is the molecular weight of the amino acid. The relation between mol fraction and S is given by:

$$x_A = \frac{S / M_A}{S / M_A + 1000 / M_w} \quad (4-50)$$

Table 4.5 lists the results for six systems, and Figure 4.3 shows the results for four systems. Since the values of the solubility are big, absolute average deviation (AAD) is used as the criterion of the correlation accuracy. As seen from the Table 4-5 and Figure 4.3, the results are satisfactory.

Table 4-5 Calculation results for solubility of amino acids

System	$\Delta s/R$	$\Delta h/R$ (K)	AAD
Alanine	0.8077	1254.9	2.81
Glycine	2.0888	1485.0	4.26
Proline	0.9102	744.9	1.48
DL-serine	4.4880	2707.6	0.44
Valine	-0.6674	1138.0	7.14
L-hydroxyproline	-0.1117	833.7	2.37

$$AAD = \frac{100}{N} \sum_{i=1}^N \frac{|S_i^{\text{exp}} - S_i^{\text{cal}}|}{S_i^{\text{exp}}}, \quad S = \text{solubility in g/kg (solvent)}$$

Figure 4.3 Correlation results for solubility of amino acids in H₂O

CHAPTER 5: CONCLUSIONS, CONTRIBUTIONS TO KNOWLEDGE AND SUGGESTIONS FOR FUTURE WORK

5.1 Conclusions

The mean ionic activity coefficients of the electrolyte and activity coefficients of the amino acids were measured at 298.15 K using the electrochemical method for the following four systems: DL-serine + NaNO₃ + Water; DL-valine + NaNO₃ + Water; DL-serine + KNO₃ + Water; glycine + KNO₃ + Water.

The activity coefficients of DL-serine in DL-serine + KNO₃ + Water system were also measured at 298.15 K using isopiestic method. The results were compared with those obtained by electrochemical method. The trends of the concentration effect on the activity coefficients from the two methods are the same. The possible error sources and advantages and disadvantages of the two methods were discussed.

The data measured for the ternary systems were correlated using the Gibbs free energy model proposed by Khoshbarchi and Vera (1996a) with the NRTL equation. Among the six binary parameters, two binary parameters were obtained from the amino acid + water system, two parameters were obtained from ternary data, and symmetrical relations were assumed for other two parameters.

A new model based on simplified perturbation theory was proposed. This model was used to correlate the measured data for the four systems. Two adjustable parameters were regressed from the ternary system, while the other parameters were obtained from binary amino acid + water systems and from the data of ions. The major advantage of the new model is the simplicity of the formulation. In addition, the amino acid molecules are characterized only by a size and an energy parameter, and no other property is required for the correlation. The results of the correlation are slightly better than those obtained from the Gibbs free energy model.

Chapter 5. CONCLUSIONS, CONTRIBUTIONS TO KNOWLEDGE AND SUGGESTIONS FOR FUTRUE WORK

The new model was also used to correlate the activity coefficients and solubilities of amino acids for some binary aqueous solution reported in literature (Fasnman, 1976). Satisfactory results were obtained.

5.2 Contributions to Knowledge

New data for mean ionic activity coefficients of electrolytes and for activity coefficients of amino acids in four ternary systems at 298.15 K were measured. The effects of the concentration of the electrolyte and the amino acid were determined.

A new model based on a simplified perturbation theory was proposed and tested with the experimental measured in this work and from literature. The model is easy to use, and no additional molecular data is needed for the calculations. The new model was applied to correlate the activity coefficients for binary (amino acid + water) and ternary (amino acid + electrolyte + water) systems and solubilities of amino acid in binary aqueous solutions.

5.3 Suggestions for Future Work

It would be of interest to measure the temperature effect on the activity coefficients, using the electrochemical and the isopiestic methods, over the temperature range from 298.15 to 323.15 K.

For comparison with the results from the electrochemical method, activity coefficients for more systems should be measured by the isopiestic method. The set up for this method can be further improved.

Based on some theoretical considerations, it could be desirable to develop a new model which can predict, with some accuracy, the activity coefficients for ternary systems using only parameters obtained from binary systems. For example, in the model based on perturbation theory, generalized mixing rules may be obtained after employing the model for many systems. New mixing rules may be necessary.

REFERENCES

- Ben-Amotz, D., Herschbarch, D. R., Estimation of Effective Diameter for Molecular Fluids, *J. Phys. Chem.*, 94, 1038, 1990.
- Bower, V. E., Robinson, R. A., Thermodynamics of the Ternary Systems: Water-Glycine-Potassium Chloride at 25 C from Vapor Pressure Measurements, *J. Res. Natl. Bureau of Standards*, 69A, 131, 1965.
- Butler, J. N., *Ionic Equilibrium: Solubility and pH Calculations*, John Wiley & Sons, New York, 1998.
- Briggs, C. C., Lilley, T. H., Rutherford, J., Woodhead, S., The Activity of Calcium Chloride in Aqueous Solutions of Some Amino Acids at 25 C, *J. Solution Chem.*, 3, 649, 1974.
- Bromley, L. A., Thermodynamic Properties of Strong Electrolytes in Aqueous Solutions, *AIChE J.*, 19, 313, 1973.
- Davies, C. W., *Ion Association*. Butterworth, London, 1962.
- Draper, N. R., Smith, H., *Applied Regression Analysis*, 2nd ed., John Wiley & Sons, Inc., New York, 1981.
- Fasman, G. D., *Handbook of Biochemistry and Molecular Biology*, 3rd ed., Physical and Chemical Data, Vol. 1 CRC Press. Cleveland, 1976.
- Haghtalab, A., *Thermodynamics of Aqueous Electrolyte Solutions*, Ph. D. Thesis, Dept. of Chem. Eng., McGill University, Montreal, 1990.

References.

Haghtalab, A., Vera, J. H., Mean Activity Coefficients in Ternary NaCl-NaNO₃-H₂O and NaBr-NaNO₃-H₂O systems at 298.15 K, *J. Chem. Eng. Data*, 36, 332, 1991.

Kelley, B. P. Lilley, T. H., Aqueous Solutions Containing Amino Acids and Peptides, V, *Chem. Soc. J., Faraday Trans. I.*, 74, 2771, 1978.

Khoshkbarchi, M. K., Vera, J. H., Measurement and Modeling of Activities of Amino Acid in Aqueous Salt Systems, *AIChE J.*, 42, 2354, 1996a.

Khoshkbarchi, M. K., Vera, J. H., Measurement of Activity Coefficients of Amino Acids in Aqueous Electrolyte Solutions: Experimental Data for the Systems H₂O + NaCl + Glycine and H₂O + NaCl + DL-Alanine at 25 °C, *IEC Res.*, 35, 2735, 1996b.

Khoshkbarchi, M. K., Vera, J. H., A Perturbed Hard-Sphere Model with Mean Spherical Approximation for the Activity Coefficients of Amino Acids in Aqueous Solutions, *IEC Res.*, 35, 4755, 1996c.

Khoshkbarchi, M. K., Vera, J. H., A Simplified Perturbed Hard-Sphere Model for the Activity Coefficients of Amino Acids and Peptides in Aqueous Solutions, *IEC Res.*, 35, 4320, 1996d.

Khoshkbarchi, M. K., Vera, J. H., Activity Coefficients of DL-Valine in Aqueous Solutions of KCl at 25 °C, Measurement with Ion Selective Electrodes and Modeling, *J. Solution Chem.*, 25, 865, 1996e.

Khoshkbarchi, M. K., Vera, J. H., Measurement and Correlation of Ion Activity Coefficients in Aqueous Single Electrolyte Solution, *AIChE J.*, 42, 249, 1996f.

References.

Khoshkbarchi, M. K., Vera, J. H., Effect of NaCl and KCl on the Solubility of Amino Acids in Aqueous Solutions at 298.2 K: Measurements and Modeling, *IEC Res.*, **36**, 2445, 1997.

Khoshkbarchi, M. K., Soto-Campos, A. M., Vera, J. H., Interaction of DL-serine and L-serine with NaCl and KCl in Aqueous Solutions, *J. Solution Chemistry*, **26**, 914, 1997.

Khoshkbarchi, M. K., Vera, J. H., A Theoretical Improved Perturbation Model for Activity Coefficients of Amino Acids and Peptides in Aqueous Solutions, *IEC Res.*, **37**, 3052, 1998.

Lewis, G. N., Randall, M., *Thermodynamics*, McGraw-Hill, New York, 1961.

Lin, D.-D., Mei, L.-H., Zhu, Z.-Q., Han, Z.-X., An Improved Isopiestic Method for Measurement of water Activities in Aqueous Polymer and Salt Solution, *Fluid Phase Equilibria*, **118**, 241, 1996.

Maitland, G. C., Rigby, M., Smith, E. B., Wakeham, W. A., *Intermolecular Forces*, Clarendon Press, Oxford, UK, 1981.

Mansoori, G. A., Carnahan, N. F., Starling, K. E., Leland, T. W., Equilibrium Thermodynamic Properties of Mixtures of Hard Spheres, *J. Chem. Phys.*, **54**, 1523, 1971.

Ochs, L. R., Kabiri-Badr, M., Cabezas, H. Jr., An Improved Isopiestic Method to Determine Activities in Multicomponent Mixtures, *AIChE J.*, **36**, 1908, 1990.

Park, H., Englezos, P., Osmotic Coefficient Data for Na_2SiO_3 and $\text{Na}_2\text{SiO}_3\text{-NaOH}$ by an Isopiestic Method and Modeling Using Pitzer's Model, *Fluid Phase Equilibria*, **153**, 87, 1998.

References.

Phang, S., Steel, B. J., Activity Coefficients from e.m.f. measurements using cation-responsive glass electrode. NaCl + Glycine + Water at 273.13, 283.15, 298.15, and 323.15 K, *J. Chem. Thermodynamics*, 6, 537, 1974.

Pradhan, A. A., Vera, J. H., Effect of Acids and Bases on the Solubility of Amino Acids, *Fluid Phase Equilibria*, 152, 121, 1998.

Robinson, R. A., Stokes, R. H., *Electrolyte Solutions*, 2nd ed., Butterworth Scientific Publications, London, 1959.

Rodriguez-Raposo, R., Fernandez-Merida, Estes, M. A., Activity Coefficients in (Electrolyte + Amino Acid) (Aq). *J. Chem. Thermodynamics*, 26, 1121, 1994.

Scatchard, G., Hamer, W. J., Wood, S. E., Isotonic Solutions 1, *J. Am. Chem. Soc.*, 60, 3061, 1938.

Schrier, E. E., Robinson, R. A., A Study of Free Energy Relationships in Some Amino Acid-Sodium-Water Systems, *J. Biol. Chem.*, 246, 2870, 1971.

Schrier, E. E., Robinson, R. A., Free Energy Relationships in Aqueous Amino Acid and Peptide Solutions Containing Sodium Chloride, *J. Solution Chem.*, 3, 493, 1974.

Silverberg, M., Activity Coefficients of DL-serine in Aqueous KNO₃, Project report (302-495B), 1998, Chem. Eng. Dept., McGill University.

Sinclair, D. A., A Simple Method for Accurate Determination of Vapor Pressures of Solutions, *J. Phys. Chem.*, 37, 495, 1933.

References.

Skoog, D. A., Leary, J. J., *Principles of Instrumental Analysis*, Fort worth, Saunders College Pub., 1992.

Soto-Campos, A. M., Khoshkbarchi, M. K., Vera, J. H., Interaction of DL-threonine with NaCl and NaNO₃ in Aqueous Solutions: e.m.f. Measurements with Ion-selective Electrodes, *J. Chem. Thermodynamics*, 29, 609, 1997a.

Soto-Campos, A. M., Khoshkbarchi, M. K., Vera, J. H., Activity Coefficients of the Electrolyte and the Amino Acid in Water + NaNO₃ + Glycine and Water + NaCl + DL-methionine Systems at 298.15 K, *Biophysical Chemistry*, 67, 97, 1997b.

Soto-Campos, A. M., Khoshkbarchi, M. K., Vera, J. H., Effect of the Anion and the Cation of an Electrolyte on the Activity Coefficient of DL-alanine in Aqueous Solutions, *Fluid Phase Equilibria*, 142, 193, 1998.

Taghikhani, V., Khoshkbarchi, M. K., Vera, J. H., On the Expression for the Chemical Potential in Mixtures of Hard Spheres. *Fluid Phase Equilibria*, 165, 141, 1999.

Tiepel, E. W., Gubbins, K. E., Thermodynamic Properties of Gases Dissolved in Electrolyte Solutions, *Ind. Eng. Chem. Fundam.*, 12, 18, 1973.

Thiessen, D. B., Wilson, A. J., An Isopiestic Method for Measurement of Electrolyte Activity Coefficients, *AIChE J.*, 33, 1926, 1987.

Zaytsev, I. D., Aseyev, M. A., *Properties of Aqueous Solutions of Electrolytes*, CRC Press, Boca Raton, 1992.

APPENDICES

A ΔE Data by Electrochemical Method

Table A.1 ΔE data for $\text{NaNO}_3 + \text{DL-serine} + \text{H}_2\text{O}$ system

m $\text{NaNO}_3 \Rightarrow$	0.1	0.3	0.5	0.7	1.0
m DL-serine $\Downarrow$	ΔE				
0	171.3	221.6	243.6	258.1	273.5
0.05	170.7	221.1	243.2	257.9	273.3
0.1	170.2	220.8	242.9	257.6	273.1
0.15	169.7	220.4	242.6	257.4	272.8
0.2	169.3	220.0	242.3	257.1	272.6
0.25	168.9	219.7	242.0	256.8	272.3
0.3	168.4	219.3	241.6	256.5	272.1
0.35	168.0	218.9	241.4	256.3	271.9
0.4	167.6	218.6	241.1	256.0	271.7

Table A.2 ΔE data for $\text{NaNO}_3 + \text{DL-valine} + \text{H}_2\text{O}$ system

m $\text{KNO}_3 \Rightarrow$	0.1	0.3	0.5	0.7	1.0
m DL-valine $\Downarrow$	ΔE				
0	174.4	225.4	248.2	262.6	278.0
0.1	174.3	225.3	248.3	262.8	278.2
0.2	174.0	225.2	248.2	262.9	278.4
0.3	173.8	225.1	248.2	262.8	278.5
0.4	173.5	224.8	248.1	262.8	278.6
0.5	173.2	224.7	248.2	262.8	278.6

Appendices

Table A.3 ΔE data for $\text{KNO}_3 + \text{DL-serine} + \text{H}_2\text{O}$ system

m $\text{KNO}_3 \Rightarrow$	0.1	0.3	0.5	0.7
m DL-serine $\Downarrow$	ΔE			
0	-32.6	10.0	26.3	35.0
0.01	-32.8	9.9	26.3	35.0
0.03	-32.9	9.8	26.2	35.0
0.05	-33.0	9.7	26.2	34.9
0.1	-33.5	9.4	26.0	34.7
0.15	-33.9	9.2	25.8	34.6
0.2	-34.5	8.9	25.6	34.5
0.25	-34.9	8.6	25.4	34.3
0.3	-35.3	8.3	25.2	34.3
0.35	-35.7	8.0	25.0	34.1
0.4	-36.1	7.7	24.9	33.9

Table A.4 ΔE data for $\text{KNO}_3 + \text{glycine} + \text{H}_2\text{O}$ system

m $\text{KNO}_3 \Rightarrow$	0.1	0.3	0.5	0.7
m glycine $\Downarrow$	ΔE			
0	-32.7	9.6	25.1	34.2
0.1	-33.6	8.9	24.6	33.9
0.3	-35.3	7.7	23.6	33.1
0.7	-38.4	5.5	22.1	31.7
1.0	-40.5	4.0	20.8	30.7
1.3	-42.5	2.5	19.7	29.9
1.6	-44.2	1.3	18.7	29.2
2.0	-46.3	-0.3	17.4	28.0
2.3	-47.8	-1.2	16.7	27.3
2.6	-49.0	-2.2	16.0	26.7

B. Derivation of Eq.(4-15)

The intermolecular force is composed of several parts: dispersion force, dipole-dipole interaction force, and induction force, etc. The dispersion can be expressed by the Lennard-Jones potential. The dipole-dipole interaction potential and induction potential for a pure substance (here in a binary amino acid + water system, the amino acid acts like a 'pure' substance in a background of continuum) are given by eqs.(B-1) and (B-2), respectively (Maitland et al., 1981).

$$u_{DD}(r) = -\frac{2}{3} \frac{D^4}{(4\pi\epsilon_0\epsilon_r)^2 kT r^6} \quad (\text{B-1})$$

$$u_{ind,DD}(r) = -\frac{2\bar{\alpha} D^2}{(4\pi\epsilon_0\epsilon_r)^2 r^6} \quad (\text{B-2})$$

where $\bar{\alpha}$ is the polarizability. Combining the all interaction potentials and inserting the potential and eq.(4-13) into eq.(4-12), the perturbed chemical potential is given by:

$$\begin{aligned} \mu_i^{Per} &= 4\pi\rho \int_{\sigma}^{\infty} u(r) r^2 dr \\ &= 4\pi\rho \int_{\sigma}^{\infty} \left\{ 4\epsilon \left[\left(\frac{\sigma}{r} \right)^{12} - \left(\frac{\sigma}{r} \right)^6 \right] - \frac{2}{(4\pi\epsilon_0\epsilon_r)^2} \left(\frac{D^4}{3kT r^6} + \frac{D^2\bar{\alpha}}{r^6} \right) \right\} r^2 dr \end{aligned} \quad (\text{B-3})$$

After integration and rearrangement, and using the relation,

$$\mu_i^{Per} = kT \ln \gamma_i^{Per} \quad (\text{B-4})$$

we have

$$\ln \gamma_i^{Per} = -\frac{4\pi\rho\sigma^3\epsilon}{kT} B \quad (\text{B-5})$$

Appendices

where B is defined as

$$B = \frac{8}{9} + \frac{2}{3(4\pi\epsilon_0\epsilon_r)^2} \left(\frac{D^4}{3kT} + D^2\bar{\alpha} \right) \frac{1}{\sigma^6\epsilon} \quad (\text{B-6})$$

Strictly, B is a function of D , $\bar{\alpha}$ and T . In this work, we found it suffices to assume $B = 0.2$ in our correlation. This significantly simplified the model.

C. Derivation of the Perturbation Term for Mixtures

The perturbed chemical potential is given by eq.(4-26). As in the 'pure' amino acid case, a step function is used for the radial distribution function. For illustration purpose, only Lennard-Jones potential is considered here (since a constant (0.2) will be finally adopted as the case discussed in Appendix B):

$$u_{ij}^L(r) = 4\epsilon_{ij} \left(\frac{\sigma_{ij}^{12}}{r^{12}} - \frac{\sigma_{ij}^6}{r^6} \right) \quad (\text{C-1})$$

In our application, the solution is considered as 'binary', after integration the double summation term is given by:

$$[\text{double sum}] = -\frac{8}{9} \left(\rho_A^2 \epsilon_{AA} \sigma_{AA}^3 + 2\rho_A \rho_S \epsilon_{AS} \sigma_{AS}^3 + \rho_S^2 \epsilon_{SS} \sigma_{SS}^3 \right) \quad (\text{C-2})$$

Assuming (1) symmetrical relations: $\epsilon_{AS} = \epsilon_{SA}$ and $\sigma_{AS} = \sigma_{SA}$, (2) the molecular energy parameters are independent of density, while the size parameters are the functions of density given by eq.(4-27), after rearrangement, the activity coefficients of amino acid and electrolyte are given by following equations:

$$\ln \gamma_A^{per} = -\frac{B\pi}{kT} \left[\rho_A \epsilon_{AA} \sigma_{AA}^3 + \rho_S \epsilon_{AS} \sigma_{AS}^3 + 3\rho_A \rho_S \epsilon_{AS} \sigma_{AS}^2 \left(\frac{\partial \sigma_{AS}}{\partial \rho_A} \right) \right] \quad (\text{C-4})$$

Appendices

$$\ln \gamma_s^{per} = -\frac{B\pi}{kT} \left[\rho_s \epsilon_{ss} \sigma_{ss}^3 + \rho_A \epsilon_{AS} \sigma_{AS}^3 + 3\rho_A \rho_s \epsilon_{AS} \sigma_{AS}^2 \left(\frac{\partial \sigma_{AS}}{\partial \rho_s} \right) \right] \quad (C-5)$$

where the constant B is taken as 0.2 to be consistent with the 'pure' amino acid case.

D. The Cross Differential Relation for the Perturbation Theory

According to thermodynamics, for the activity coefficients of the amino acid + electrolyte + water system, the following cross differential relation should be satisfied

$$\left(\frac{\partial \ln \gamma_A}{\partial m_s} \right)_{T, m_1} = v \left(\frac{\partial \ln \gamma_s}{\partial m_A} \right)_{T, m_s} \quad (D-1)$$

or using the definition of number density,

$$\left(\frac{\partial \ln \gamma_A}{\partial \rho_s} \right)_{T, \rho_1} = v \left(\frac{\partial \ln \gamma_s}{\partial \rho_A} \right)_{T, \rho_s} \quad (D-2)$$

Combining eqs.(4-4) and (4-5) with eq.(D-2), we have

$$\left(\frac{\partial \ln \gamma_A^{HS}}{\partial \rho_s} \right)_{T, \rho_1} = \left(\frac{\partial \ln \gamma_s^{HS}}{\partial \rho_A} \right)_{T, \rho_s} \quad (D-3)$$

and

$$\left(\frac{\partial \ln \gamma_A^{per}}{\partial \rho_s} \right)_{T, \rho_1} = \left(\frac{\partial \ln \gamma_s^{per}}{\partial \rho_A} \right)_{T, \rho_s} \quad (D-4)$$

Eq.(D-3) is automatically satisfied according to the derivation of the activity coefficients of hard sphere mixture. Eq.(D-4) is satisfied according to the derivation of the perturbation activity coefficient and the mixing rules eqs.(4-27) and (4-28). In fact, it is easy to show that

Appendices

$$\left(\frac{\partial \ln \gamma_A^{Per}}{\partial \rho_S} \right)_{\rho_A} = \left(\frac{\partial \ln \gamma_S^{Per}}{\partial \rho_A} \right)_{\rho_S} =$$

$$-\frac{B\pi}{kT} \varepsilon_{AS} \sigma_{AS}^3 \left[1 + \frac{9}{2} \rho_A \rho_S \delta^{(1)} \frac{\sigma_{AA} + \sigma_{SS}}{\sigma_{AS}} + \frac{3}{2} (\rho_A \rho_S \delta^{(1)})^2 \left(\frac{\sigma_{AA} + \sigma_{SS}}{\sigma_{AS}} \right)^2 \right]$$

(D-5)

Therefore, the equations used in this work satisfy the cross differential relation.