ON THE PERFECT ORDERABILITY OF UNIONS OF TWO GRAPHS

A THESIS

SUBMITTED TO THE FACULTY OF GRADUATE STUDIES AND RESEARCH

IN PARTIAL FULFILLMENT OF THE REQUIREMENTS

FOR THE DEGREE OF

MASTER OF SCIENCE

IN

COMPUTER SCIENCE

FACULTY OF ARTS AND SCIENCE

LAKEHEAD UNIVERSITY

By

Xiaodan Tu

Thunder Bay, Ontario

August 1996

© Copyright 1996: Xiaodan Tu

National Library
of Canada

Acquisitions and
Bibliographic Services

395 Wellington Street
Ottawa ON K1A 0N4
Canada

Bibliothèque nationale
du Canada

Acquisitions et
services bibliographiques

395, rue Wellington
Ottawa ON K1A 0N4
Canada

Your file Votre référence

Our file Notre référence

The author has granted a non-exclusive licence allowing the National Library of Canada to reproduce, loan, distribute or sell copies of this thesis in microform, paper or electronic formats.

The author retains ownership of the copyright in this thesis. Neither the thesis nor substantial extracts from it may be printed or otherwise reproduced without the author's permission.

L'auteur a accordé une licence non exclusive permettant à la Bibliothèque nationale du Canada de reproduire, prêter, distribuer ou vendre des copies de cette thèse sous la forme de microfiche/film, de reproduction sur papier ou sur format électronique.

L'auteur conserve la propriété du droit d'auteur qui protège cette thèse. Ni la thèse ni des extraits substantiels de celle-ci ne doivent être imprimés ou autrement reproduits sans son autorisation.

0-612-33459-7

Canada

Table of Contents

Table of Contents	i
List of Tables	iii
List of Figures	iv
Chapter 1 INTRODUCTION	1
1.1 Graph Coloring Problem	1
1.2 Applications	2
1.3 Overview of the Thesis	3
Chapter 2 GRAPH COLORING TECHNIQUES AND PERFECTLY ORDERABLE GRAPHS	4
2.1 NP-Completeness	4
2.2 Maximal Independent Set Approach	5
2.3 Sequential Approach	7
2.4 Perfectly Orderable Graph	9
2.5 Definition and Theorem	9
2.6 Recognizing Perfectly Orderable Graphs is NP-Complete	11
2.7 Orientations and Perfect Orderability	12
2.8 Some Known Subclasses of Perfectly Orderable Graphs	13

2.8.1	Comparability Graphs	13
2.8.2	Triangulated Graphs and Cotriangulated Graphs	14
2.8.3	P_4 -Indifference Graphs	16
2.8.4	P_4 -Comparability Graphs	16
2.8.5	P_4 -Simplicial Graphs	17
2.8.6	Raspail Graphs	17
2.8.7	Welsh-Powell Perfect Graphs and Matula Perfect Graphs . . .	18
2.8.8	Graphs with Dilworth Number at Most Three	19
2.8.9	The Union of Two Threshold Graphs	19
2.8.10	Intersection of Two Threshold Graphs	22
2.8.11	D-Graphs	23
2.8.12	Brittle Graphs	23

Chapter 3	INVESTIGATION OF THE PERFECT ORDERABILITY OF THE UNION OF TWO GRAPHS	25
3.1	Motivation	25
3.1.1	The List of Problems	26
3.1.2	Explanation of the List	26
3.2	The Perfect Orderability of the Union of Two Graphs in Three Cases	30
3.2.1	Some Minimally Non-perfectly Orderable Graphs	30
3.2.2	Comparison with Known Classes of Perfectly Orderable Graphs	36
3.2.3	Conjectures	40
3.3	Proof of the Theorems	41
3.4	Discussion of the Proof of Theorem 3.3	50

List of Tables

2.1	Six Possible Subclasses of Perfectly Orderable Graphs.	16
3.1	The List of the Union of Two Graphs.	27

List of Figures

2.1		11
2.2	An obstruction.	12
2.3		13
2.4	Transitive orientation.	14
2.5	Semi-transitive orientation.	14
2.6	Simplicial orientation.	15
2.7		16
2.8	A house.	17
2.9		20
2.10		21
2.11		21
2.12		21
3.1		26
3.2	C_5	28
3.3	C_5	29
3.4	C_5	29
3.5	Possible G_1	31
3.6		32
3.7	An anti-hole.	32

3.8	Example 1.	33
3.9	Example 2.	34
3.10	Example 3.	35
3.11	Example 1.	36
3.12		38
3.13		38
3.14	A non-D-graph.	39
3.15		40
3.16		42
3.17		43
3.18		43
3.19		44
3.20		45
3.21		46
3.22		47
3.23		48
3.24		48
3.25	Two possible P_4	49
3.26		49
3.27	A counter-example to the rule.	51
3.28	A counter-example to the question.	52

ACKNOWLEDGEMENT

I would like to express my sincere thanks to my thesis supervisor, Dr. C.T. Hoàng, for his patience, constant encouragement, and guidance in this study. I will remain indebted to him for introducing me to scientific research.

I thank the two anonymous referees for pointing out several mistakes in the first version of this thesis. I would also like to thank others who have helped me in many different ways. The Department of Computer Science gave me academic instructions and provided convenient facilities. My classmates and friends, Jian Wang and Jun Wang, offered me help and advice. The Office of Graduate Studies and Research and my supervisor, Dr. Hoàng, provided financial assistance during the period of my study. My parents and sister gave me special encouragement and care all the time. Without these help, I would find it difficult to undertake and complete this study. To all of them I express my gratitude.

Abstract

A graph G is *perfectly orderable* if it admits an order $<$ on its vertices such that the sequential coloring algorithm delivers an optimum coloring on each induced subgraph $(H, <)$ of $(G, <)$. A graph is a *threshold graph* if it contains no P_4 , $2K_2$, or C_4 as induced subgraph. A theorem of Chvátal, Hoàng, Mahadev and de Werra states that a graph is perfectly orderable if it can be written as the union of two threshold graphs. In this thesis, we investigate possible generalizations of the above theorem. We conjecture that if G is the union of two graphs G_1 and G_2 then G is perfectly orderable whenever (i) G_1 and G_2 are both P_4 -free and $2K_2$ -free, or (ii) G_1 is P_4 -free, $2K_2$ -free and G_2 is P_4 -free, C_4 -free. We show that the complement of the chordless cycle with at least five vertices cannot be a counter-example to our conjecture and we prove, jointly with Hoàng, a special case of (i): if G_1 and G_2 are two edge-disjoint graphs that are P_4 -free and $2K_2$ -free then the union of G_1 and G_2 is perfectly orderable.

Chapter 1

INTRODUCTION

In this chapter, a brief introduction to the graph coloring problem and its application is given, followed by an overview of the thesis.

1.1 Graph Coloring Problem

Graph coloring problem is to color the vertices of a graph using a minimum number of colors, subject to the restriction that no two adjacent vertices get the same color. Ever since it was originally formulated in the last century, finding an efficient way to optimally color a graph has attracted the interests of mathematicians. Although many significant results have been derived in this field, it is still left to be one of the most intractable problems in discrete mathematics.

We now give the definition of the graph coloring problem in the term of graph theory:

Definition 1.1 *A graph G is said to be r -colorable if its vertices can be colored with r colors in such a way so that no two adjacent vertices are of the same color. The smallest number r for which the graph is r -colorable is called the chromatic number $\chi(G)$ of the graph and finding this number is referred to as the coloring problem.*

Here, G is assumed to be a simple graph, that is, an undirected graph with neither loops nor multiple edges.

1.2 Applications

The earliest application of graph coloring can be traced back to the last century when it was originally formulated. The cartographers asked for a minimum number of colors to color a political map in such a way that no two neighbouring countries get the same color. In this example, different countries can be regarded as the set of vertices V in G , and the two vertices of V are adjacent if and only if the two countries they represent are neighbours. They predicted that four colors always suffice. This is known as the *Four Color Problem*, which was solved by Appel and Haken ([1], [2]) in 1977.

A contemporary application of the coloring problem is the scheduling and loading problem, as illustrated in the following example.

A computer program stores the values of its variables in memory. For arithmetic computations, the values must be entered in "registers". Registers are expensive, so we want to use them efficiently. If two variables are not used at the same time, we can allocate them to the same register. For each variable, we compute the first and last time it is used. A variable is *active* during the interval between these times. We define a graph with the variables as vertices, in which two vertices are adjacent if they are active at a common time. The number of registers needed is the optimum coloring of the corresponding graph. Such a graph is called an *interval graph*:

Definition 1.2 *Given any family of intervals, we can define a graph whose vertices are the intervals, with vertices adjacent when the intervals intersect. A graph formed in this way is an interval graph, and the family of intervals is an interval representation of the graph.*

Interval graphs belong to the class of triangulated graphs, the optimum coloring of which can be obtained in polynomial time ([9]). We will discuss triangulated graphs in more detail later.

1.3 Overview of the Thesis

This thesis is concerned with the perfect orderability of the union of two graphs. The concept of perfectly orderable graph was developed with the motivation to solve the graph coloring problem. After a brief introduction to the graph coloring problem and its application, in chapter 2, we shall give a brief survey of two graph coloring techniques: maximal independent set and sequential approaches. We shall introduce "perfectly ordered graphs" which are the ordered graphs for which the sequential algorithm delivers an optimum coloring. Chvátal, Hoàng, Mahadev and de Werra proved that a graph is perfectly orderable if it is the union of two threshold graphs. In chapter 3, we shall investigate possible generalizations of this theorem. In particular, we conjecture that if G is the union of two graphs G_1 and G_2 then G is perfectly orderable whenever (i) G_1 and G_2 are both P_4 -free and $2K_2$ -free, or (ii) G_1 is P_4 -free, $2K_2$ -free and G_2 is P_4 -free, C_4 -free. We show that the complement of the chordless cycle with at least five vertices cannot be a counter-example to our conjecture and we prove, jointly with Hoàng, a special case of (i): if G_1 and G_2 are two edge-disjoint graphs that are P_4 -free and $2K_2$ -free then the union of G_1 and G_2 is perfectly orderable.

Chapter 2

GRAPH COLORING TECHNIQUES AND PERFECTLY ORDERABLE GRAPHS

In this chapter, a further discussion of the graph coloring is given. Despite the NP-complete property of this problem, we introduce two basic approaches to color a graph: maximal independent set and sequential, followed by a more detailed addressing on the concept and study of perfectly orderable graphs.

2.1 NP-Completeness

The coloring problem was proved to belong to the class of NP-complete problems ([7]), and worse than that, a polynomial approximation algorithm with a constant error ratio cannot exist unless $P = NP$ ([16]).

As a result of its complexity, although the number of papers on the coloring problem exceeds that on any other graph problem, no formula has been found for the chromatic number of an arbitrary graph and we must thus be satisfied with bound estimates. Here, we are going to introduce only two simple bounds:

$$\text{i) lower bound : } \omega(G) \leq \chi(G) \quad (2.1)$$

$\omega(G)$ is the number of vertices in the largest clique of G . Since all vertices of any clique of G must have different colors, this lower bound is obvious.

$$\text{ii) upper bound : } \chi(G) \leq \Delta(G) + 1 \quad (2.2)$$

$\Delta(G)$ denotes the maximum degree of a vertex of G . This inequality follows from the observation that if $\Delta(G) + 1$ colors are available, then at each vertex v of the graph G at least one of the colors can be used, since at most $\Delta(G)$ colors are used to color the neighbours of v . Brooks ([18]) proved further that there are only two classes of graphs for which the upper-bound holds with equality: odd cycles and complete graphs.

Despite the great difficulty, many efforts have been given to tackle the coloring problem. There are basically two approaches: independent set and sequential.

2.2 Maximal Independent Set Approach

A k -coloring of G is a partition of the set of vertices of G into k independent sets $V_1, V_2, \dots, V_k$ such that $V_i \cap V_j = \emptyset$ for $i \neq j, i, j = 1, 2, \dots, k$, and $\bigcup_{i=1}^k V_i = V$. Such a partition is called a *k -coloring partition* of V . Thus, the coloring problem is equivalent to finding a minimum k in a k -coloring partition of G .

Let $V_1, V_2, \dots, V_k$ be a k -coloring partition of a k -colorable graph G . Then we can construct a l -coloring partition $V'_1, V'_2, \dots, V'_l$ of G such that V'_1 is a maximal (in the sense of set-inclusion) independent set of G , and $l \leq k$ in the following way: First set $V'_1 = V_1$, and for each vertex x in $V_2 \cup V_3 \dots \cup V_k$, we put x in V'_1 if and only if x has

no neighbour in V_1' . Then we set $V_i' = V_i - \{x | x \in V_1'\}$ for $2 \leq i \leq k$. It follows that there exists a maximal independent subset U of the vertices of G such that

$$\chi(G) = \chi(G_{V-U}) + 1 \quad (2.3)$$

(G_{V-U} is the subgraph of G induced by $V - U$.)

There is a finite number of maximal independent sets W in G . minimizing over all such subsets, we obtain

$$\chi(G) = \min_{W \subseteq U} \chi(G_{V-W}) + 1 \quad (2.4)$$

Equation 2.4 is the basis for the following algorithm (Maximal Independent Set approach, or MIS) for computing the chromatic number:

Procedure MIS(G, k);

Input: a non-empty graph G .

output: chromatic number k of G .

begin {procedure}

 if $G = \emptyset$ then $k := 0$

 else

 begin

$k := |V(G)|$;

 for each maximal independent set S of G do

 begin

 MIS($G - S, l$);

 if $(l + 1) < k$ then $k := l + 1$;

 end;

 end;

end; {procedure}

It is easy to see that procedure MIS can be modified to produce a k -coloring of a k -chromatic graph. The complexity of this method is $O(mn2.445^n)$ (see [19], as usual n and m denote the number of vertices and edges respectively).

2.3 Sequential Approach

Coloring function is often used when discussing the sequential approach for coloring a graph.

Definition 2.1 *A function f determines a k -coloring of G , if*

$$f : V \rightarrow \{1, 2, \dots, k\} \quad (2.5)$$

with $f(i) \neq f(j)$ for all $(i, j) \in E$. A function that defines a k -coloring is called the k -coloring function.

A *sequential approach* can be stated as the following greedy algorithm:

Algorithm (Greedy coloring) The *greedy coloring* with respect to a vertex ordering $v_1, \dots, v_n$ of $V(G)$ is obtained by coloring vertices in the order $v_1, \dots, v_n$, assigning to v_i the smallest-indexed color not already used on its lower-indexed neighbours.

From the above sequential approach, it is easy to determine an upper bound $u_s(G; v_1, v_2, \dots, v_n)$ for the number of colors $\chi_s(G)$ used by the sequential algorithm applied to G and the ordering of its vertices $v_1, v_2, \dots, v_n$. Every vertex v_i can be colored by color i , therefore $f(v_i) \leq i$. On the other hand, at least one of the first $\deg(v_i) + 1$ colors can be assigned to v_i . Hence,

$$f(v_i) \leq \min\{i, \deg(v_i) + 1\} \quad (2.6)$$

for every $i = 1, 2, \dots, n$, and thus

$$\chi_s(G) \leq u_s(G; v_1, v_2, \dots, v_n) = \max_{i \leq n} \min\{i, \deg(v_i) + 1\} \quad (2.7)$$

This inequality is obtained without any assumption about the ordering of vertices of G .

It is obvious that in a sequential approach, the key point is to order the vertices of G , because it exclusively decides the behaviour of the greedy algorithm.

Welsh and Powell ([6]) gave the first version of sequential method by ordering the vertices according to nonincreasing degree, $\deg(v_1) \geq \deg(v_2) \geq \dots \geq \deg(v_n)$. Such an ordering is called the largest-first ordering, or LF.

A closer inspection of the algorithm and the proof of the inequality(2.7) reveals that for a given ordering $v_1, v_2, \dots, v_n$ of the vertices of a graph G , instead of $f(v_i) \leq 1 + \deg(v_i)$ we have in fact $f(v_i) \leq 1 + \deg_i(v_i)$, where $\deg_i(v_i)$ denotes the degree of vertex v_i in the subgraph of G induced by $v_1, v_2, \dots, v_i$. Therefore, the algorithm never requires more than $\max\{1 + \deg_i(v_i) : 1 \leq i \leq n\}$ colors; hence

$$\chi_s(G) \leq u'_s(G; v_1, v_2, \dots, v_n) = 1 + \max_{1 \leq i \leq n} \deg_i(v_i) \quad (2.8)$$

The following procedure finds a vertex ordering which minimizes $u'_s(G; v_1, v_2, \dots, v_n)$:

1. v_n is a minimum degree vertex of G .
2. For $i = n - 1, n - 2, \dots, 2, 1$, v_i is a minimum degree vertex in the subgraph of G induced by $V - \{v_n, v_{n-1}, \dots, v_{i+1}\}$.

Such an ordering is called smallest-last SL.

Both LF and SL algorithms may improve the upper bound (2.2) substantially. But neither of them can guarantee an optimum coloring of an arbitrary graph G . It

is natural for one to ask such a question: for which kind of ordered graphs does the greedy algorithm produce an optimum coloring? (An ordered graph is a graph with a given total order $<$ on its vertices.) This is what motivated V.Chvátal to propose the concept of perfectly orderable graph.

2.4 Perfectly Orderable Graph

In this section, we introduce the concept of perfectly orderable graph, and the well-known theorem developed by Chvátal that reveals the equivalent nature of perfectly orderability and admissible order. Knowing that to recognize perfectly orderable graphs is NP -complete, we introduce some subclasses of perfectly orderable graphs, many of which can be recognized in polynomial time.

2.5 Definition and Theorem

Definition 2.2 (Chvátal [3]) *An obstruction in an ordered graph $(G, <)$ is a set of four vertices a, b, c, d with edges ab, bc, cd (and no other edges) and $a < b, d < c$. A linear order on the set of vertices of a graph will be called i) admissible if it creates no obstruction and ii) perfect if, for each induced ordered subgraph H , the greedy algorithm produces an optimum coloring of H . A graph will be called perfectly orderable if it admits a perfect order.*

Chvátal also revealed the nature between perfect order and admissible order by proving the following theorem, which becomes the criterion for identifying perfectly orderable graphs:

Theorem 2.1 (Chvátal [3]) *A linear order of the set of vertices of a graph is perfect if and only if it is admissible.*

Proof: Since the class of graphs having obstruction-free orderings is hereditary (the inherited ordering for an induced subgraph is obstruction-free), it suffices to show that an obstruction-free ordering L gives a greedy coloring of G that is optimum. If the greedy coloring uses k colors for the ordering L , then optimality can be established by showing that G has a k -clique (a clique on k vertices). To show that such a clique does exist, we introduce the following lemma:

Lemma 2.1 *Suppose that G has a clique Q and a stable set S disjoint from Q , and suppose that for each vertex $w \in Q$ there is a vertex $p(w) \in S$ such that $p(w)$ and w are adjacent. If L is an obstruction-free ordering of G such that $p(w) < w$ for all $w \in Q$, then some $p(w) \in S$ is adjacent to all of Q .*

Proof: By induction on $|Q|$. The lemma holds trivially for $|Q| = 1$, so we may assume $|Q| > 1$. For each $w \in Q$, the graph $Q - w$ satisfies the hypotheses using the clique $Q - w$ and the stable set $\{p(u) : u \in Q - w\}$. By the induction hypothesis there is a vertex $w^* \in Q - w$ such that $p(w^*)$ is adjacent to all $Q - w$. We may assume that $p(w^*)$ is not adjacent to w for every $w \in Q$ (for otherwise, $p(w^*)$ is adjacent to all of Q and we are done). This assigns a unique w^* to every w , since $p(w^*)$ is nonadjacent only to w among Q . So setting $\sigma(w) = w^*$ defines a bijection σ on the vertices of Q .

Let v be the least vertex of Q in L . Let $b, c \in Q$ be the vertices such that $b^* = v$ and $c^* = b$. Let $a = p(b)$ and $d = p(v)$. Because $p(w^*)$ is not adjacent to w and $p(w) < w$, we have that d is not adjacent to b , a is not adjacent to c , and $d < v$ and $a < b$. Since a, d belong to the stable set S , a is not adjacent to d either (see the Figure 2.1 below. An oriented edge of the form $x \rightarrow y$ means $x < y$ and $xy \in E$.)

Because $d = p(b^*)$, the only vertex of Q nonadjacent to d is b , which implies that c is adjacent to d . Since $d = p(v) < v < c$ in L , the edge dc is oriented as $d \rightarrow c$, which implies that a, b, c and d induce an obstruction, a contradiction. $\square$

Now we continue to prove the Theorem 2.1.

Figure 2.1

Let $f : V(G) \rightarrow \{1, 2, \dots, k\}$ be the coloring generated by the greedy algorithm with this ordering. Let i be the smallest integer such that G has a clique consisting of vertices $w_{i+1}, \dots, w_k$ such that $f(w_j) = j$. Since f uses k on some vertex, such a clique exists. If $i = 0$, then G has a k -clique. Suppose $i > 0$. For each w_j there is a vertex $p(w_j)$ such that $p(w_j) < w_j$ in L and $f(p(w_j)) = i$; otherwise the greedy coloring would have used a lower color on w_j . Since the vertices in $S = \{p(w_{i+1}), \dots, p(w_k)\}$ all have color i , S is a stable set. Hence, the conditions of the Lemma 2.1 are satisfied, and there is a vertex of S that can be added to the clique and called w_i , which contradicts the minimality of i . $\square$

Perfectly orderable graphs generalize many well-known classes of graphs, such as comparability graphs, triangulated graphs and their complements. We shall discuss this fact later.

2.6 Recognizing Perfectly Orderable Graphs is NP-Complete

When Chvátal introduced the notion of perfectly orderable graphs, he also posed the question: how difficult is it to recognize perfectly orderable graphs? Middendorf

and Pfeiffer answered this question by proving the following theorem:

Theorem 2.2 (*Middendorf, Pfeiffer [17]*) *To decide whether a graph admits a perfect order is NP-complete. $\square$*

They proved the theorem by giving a reduction of 3SAT to the problem of deciding whether a graph admits a perfect order.

2.7 Orientations and Perfect Orderability

When studying perfectly orderable graphs, sometimes it is convenient to work with orientations instead of orders.

Definition 2.3 *An orientation U of a graph G is a directed graph obtained from G by assigning a direction to each edge of G .*

To an ordered graph $(G, <)$, there corresponds an orientation $D(G, <)$ of G such that $\overrightarrow{ab} \in D(G, <)$ if and only if $ab \in E(G)$ and $a < b$. So a graph is perfectly orderable if and only if it admits an acyclic orientation that does not contain an induced subgraph isomorphic to the Figure 2.2:

Figure 2.2: An obstruction.

Equivalently: a graph is perfectly orderable if and only if it admits an acyclic orientation in which each induced path of length three is one of the three types in Figure 2.3:

Figure 2.3

Although the class of perfectly orderable graphs has very nice properties in the sense of optimization, there is no known polynomial-time algorithm to recognize it. However, many subclasses of perfectly orderable graphs with special characteristics have been studied, many of which can be recognized in polynomial time. Next, we are going to introduce some of the known classes of perfectly orderable graphs.

2.8 Some Known Subclasses of Perfectly Orderable Graphs

2.8.1 Comparability Graphs

Definition 2.4 *A simple graph G is a comparability graph if it has a transitive orientation, which is an acyclic orientation such that if $xy, yz \in E(G)$, $x \rightarrow y$ and $y \rightarrow z$, then $xz \in E(G)$ and $x \rightarrow z$.*

From Definition 2.4, a transitive orientation implies that each induced P_3 in a comparability graph is of the types in Figure 2.4, and it is clear there is no obstruction. So comparability graphs are perfectly orderable.

Figure 2.4: Transitive orientation.

The following theorem of Ghouila-Houri ([8]) is the key to a polynomial-time algorithm to recognize comparability graphs.

Theorem 2.3 *A graph is a comparability graph if and only if it admits an orientation that contains no induced subgraph isomorphic to the graph in Figure 2.5 (a semi-transitive orientation). $\square$*

Note that in the above theorem, the orientation may contain directed cycles.

Figure 2.5: Semi-transitive orientation.

2.8.2 Triangulated Graphs and Cotriangulated Graphs

Definition 2.5 *A triangulated graph G is a graph such that every cycle of length ≥ 4 in G has a chord, that is, an edge joining two non-consecutive vertices of the cycle. It is also called chordal, rigid-circuit, monotone transitive, and perfect elimination graph.*

A triangulated graph G has the property that every induced subgraph contains a vertex whose neighbourhood induces a clique (a simplicial vertex). It follows that it admits an order $<$, such that v_i is simplicial in the subgraph induced by $H =$

$\{x|x < v_i\} \cup \{v_i\}$. Such an order is called a *simplicial order*. In a simplicial order, each P_3 is of the types shown in Figure 2.6. So it is obvious that a simplicial order is a perfect order. Testing whether a graph is triangulated is polynomially equivalent to testing whether a graph contains a simplicial vertex which can be obviously solved in polynomial time.

Figure 2.6: Simplicial orientation.

Definition 2.6 A *cotriangulated graph* is the complement of a triangulated graph.

In a triangulated graph G , the simplicial order implies that each P_4 is of the Type 1 or 2 in Figure 2.3. In the complement $\overline{G}$ of G , define an order $<_{\overline{G}}$ such that $x <_{\overline{G}} y$ if and only if $y <_G x$. We shall show that the order $<_{\overline{G}}$ contains no obstruction.

By $abcd$, we denote the P_4 with vertices a, b, c, d , edges ab, bc, cd (and no other edges). If $abcd$ is a P_4 of G , then its complement in $\overline{G}$ is $bdac$. Suppose $abcd$ is a P_4 of Type 1 in G (see Figure 2.7 (a)). Without loss of generality, we may assume that $b <_G a$, $b <_G c$, $c <_G d$. The last two relations imply that $b <_G d$, and therefore $d <_{\overline{G}} b$. Thus $bdac$ is not an obstruction in $\overline{G}$. A similar argument shows that the complement of a P_4 of Type 2 in G cannot be an obstruction (Figure 2.7 (b)).

By restricting the orientation on P_4 to each of the three types in Figure 2.3, Hoàng and Reed ([15]) introduced the following six subclasses of perfectly orderable graphs, four of which can be recognized in polynomial time.

Figure 2.7

	P_4 allowed in G			Name	Recognition Complexity
	Type 1	Type 2	Type 3		
1	✓	✓		P_4 -simplicial	Polynomial
2	✓		✓	Generalized CR	NP-Complete
3		✓	✓	One-in-one-out	Unknown
4	✓			Raspail	Polynomial
5		✓		P_4 -indifference	Polynomial
6			✓	P_4 -comparability	Polynomial

Table 2.1: Six Possible Subclasses of Perfectly Orderable Graphs.

2.8.3 P_4 -Indifference Graphs

Definition 2.7 A graph G is P_4 -indifference if there is an acyclic orientation of G in which every P_4 is of Type 2.

2.8.4 P_4 -Comparability Graphs

Definition 2.8 A graph G is P_4 -comparability if it admits an acyclic orientation in which each P_4 is of Type 3.

In a transitive orientation, every P_4 is of Type 3. Thus the class of P_4 -comparability graphs contains all comparability graphs.

Figure 2.8: A house.

2.8.5 P_4 -Simplicial Graphs

Definition 2.9 *A graph G is P_4 -simplicial if there is an acyclic orientation of G in which every P_4 of G is of Type 1 or 2.*

In a simplicial order, every P_4 is of Type 1 or 2. Thus the class of P_4 -simplicial graphs contains all triangulated graphs.

2.8.6 Raspail Graphs

Definition 2.10 *A graph is Raspail if it admits an acyclic orientation in which every P_4 is of Type 1.*

Hertz and de Werra ([11]) gave a characterization of Raspail graphs by forbidding induced subgraphs, among which is the *house*, as illustrated in Figure 2.8.

It is easy to see that in a house $abcde$, the two P_4 s $abcd$ and $cdea$ cannot be both of Type 1. We shall refer to this property of the house later in the thesis.

Hoàng and Reed ([15]) also developed polynomial-time algorithms to recognize the above four classes of perfectly orderable graphs.

Hoàng ([13]) also showed that recognizing "generalized CR" graphs is NP-complete.

It is not known whether "one-in-one-out" graphs can be recognized in polynomial time.

2.8.7 Welsh-Powell Perfect Graphs and Matula Perfect Graphs

We consider graphs G with linear orders $<$ on the vertices. Welsh and Powell choose $<$ in such a way that,

$$d_G(x) \geq d_G(y) \text{ whenever } x < y \quad (2.9)$$

with $d_G(x)$ standing for the degree of x in G ;

while Matula chooses $<$ in such a way that,

$$d_H(x) \geq d_H(y) \text{ whenever } x < y \quad (2.10)$$

and H is the subgraph of G induced by all z with $z \leq y$.

Definition 2.11 *G is called Welsh-Powell perfect if the linear order $<$ satisfying (2.9) is perfect (the greedy algorithm produces an optimum coloring for each induced subgraph of G), and Matula perfect if the linear order $<$ satisfying (2.10) is perfect.*

Chvátal, Hoàng, Mahadev and de Werra ([5]) proved theorems to characterize Welsh-Powell perfect and Matula perfect graphs by forbidding certain induced subgraphs. Again a *house* is such a forbidden subgraph for both Welsh-Powell and Matula perfect graphs. They also showed that these two classes of perfectly orderable graphs can be recognized in polynomial time, and presented an algorithm to prove the following theorem:

Theorem 2.4 *Given any graph G that is Welsh-powell perfect or Matula perfect, one can find in time $O(m + n)$ a minimum coloring and a largest clique in G . Given any graph G whose complement is Welsh-Powell perfect or Matula perfect, one can find in time $O(m + n)$ a minimum clique cover and a largest stable set in G . $\square$*

2.8.8 Graphs with Dilworth Number at Most Three

Definition 2.12 Let $N(x)$ stand for the set of all the neighbours of a vertex x in G ; we say that a vertex y dominates x if $N(x) \subseteq N(y) \cup \{y\}$. x and y are comparable if x dominates y or y dominates x , and incomparable if neither of them dominates the other. The Dilworth number of G is the largest number of pairwise incomparable vertices in G .

It is obvious that the class of graphs with Dilworth number at most three is recognizable in polynomial time. In [5], it is proved that these graphs are perfectly orderable.

2.8.9 The Union of Two Threshold Graphs

Threshold graphs were introduced to study the stable set polytope of a graph.

Definition 2.13 Let $G = (V, E)$ be a graph on n vertices. Then G is a threshold graph if there exists a linear inequality

$$\sum_{i=1}^n a_i x_i \leq a \quad (2.11)$$

with $a, a_i \in \mathbb{R}$ and $n = |V|$ such that the following holds: $S \subset V$ is a stable set of G if and only if (2.11) is satisfied by the characteristic vector $x_s = (x_1, x_2, \dots, x_n)$ of S where for all i

$$x_i = \begin{cases} 1 & \text{if } i \in S \\ 0 & \text{if } i \notin S \end{cases} \quad (2.12)$$

Theorem 2.5 (Chvátal, Hammer [4]) A necessary and sufficient condition for G to be threshold is that G does not contain $2K_2$, P_4 or C_4 as an induced subgraph (Figure 2.9.)

Figure 2.9

Chvátal, Hoàng, Mahadev and De Werra proved the following theorem:

Theorem 2.6 *If G is the union of two threshold graphs then G is perfectly orderable.*

We are going to give a new proof of the Theorem 2.6. But before that, let us first give some definitions about P_4 : Let $abcd$ denote the P_4 with vertices a, b, c, d and edges ab, bc, cd , the edges ab and cd are called *wings* of P_4 .

Proof: Let G be the union of two threshold graphs G_1 and G_2 . Consider any P_4 $v_0v_1v_2v_3$ in G (if it exists). It is clear that the two wings v_0v_1 and v_2v_3 cannot be in the same G_i ($i = 1, 2$), otherwise there will be either a $2K_2$ or a P_4 in G_i .

$\forall P_4 \in G$, we now impose a partial orientation on the edges of G :

1. direct $v_1 \rightarrow v_0$ if and only if $v_0v_1 \in G_i$ and $v_1v_2 \in G_i$,
2. direct $v_2 \rightarrow v_3$ if and only if $v_1v_2 \in G_i$ and $v_2v_3 \in G_i$.

It is obvious that such a partial orientation does not create an obstruction in G .

Claim 2.1 *Such an orientation is unique on each edge.*

Proof: Suppose there is an edge cd that receives two directions by the above orientation.

Let us assume $cd \in G_1$. We must have a P_4 $abcd$ with $ab \in G_2$, $bc \in G_1$ and another P_4 $cdef$ with $de \in G_1$, $ef \in G_2$. Since $bd \notin G_1$ and $ce \notin G_1$, $\{b, c, d, e\}$ form either a P_4 or C_4 in G_1 , a contradiction (see Figure 2.10.) $\square$

(In all of the following figures in this thesis, we use solid lines to represent G_1 and dashed lines to represent G_2 .)

Figure 2.10

Claim 2.2 *There are no two directed edges $\overrightarrow{ab}$, $\overrightarrow{bc}$ (Figure 2.11) such that $\overrightarrow{ab} \in G_1$ (respectively G_2) and $\overrightarrow{bc} \in G_2$ (respectively G_1).*

Figure 2.11

Proof: Suppose there are directed edges $\overrightarrow{ab}$, $\overrightarrow{bc}$ with $\overrightarrow{ab} \in G_1$ and $\overrightarrow{bc} \in G_2$.

Figure 2.12

Since $a \rightarrow b$ in G_1 , there must be a P_4 $deab$ with $de \in G_2$, and $ea, ab \in G_1$ (c is not identical to either d or e); similarly, since $b \rightarrow c$ in G_2 , there must be a P_4 $gfb c$

with $gf \in G_1$ and $fb, bc \in G_2$ (g may be identical to e , but f cannot be identical to either d or e). Since G_2 is a threshold graph, to have both edges ed and bc belonging to G_2 , there must be $ce \in G_2$ and $cd \in G_2$. Now consider the subgraph consisting of the three edges dc , cb and bf in G_2 . We have $db \notin E(G)$ and $cf \notin E(G)$. There will be either an induced P_4 (if $df \notin E(G_2)$) or C_4 (if $df \in E(G_2)$) in G_2 , a contradiction (see Figure 2.12.) $\square$

Claim 2.3 *The partial orientation creates no directed cycle in G .*

Proof: We note that for any directed edge $\overrightarrow{ab} \in G_i$, a strictly dominates b in G_i (a dominates b , but b does not dominate a). Therefore, there will be no directed cycle in the same G_i . Furthermore, Claim 2.2 guarantees that there is no directed cycle consisting of edges in both G_1 and G_2 . $\square$

Now we can easily extend this acyclic partial orientation to a linear order $<$ on G by the following procedure:

1. $i \leftarrow 1, H \leftarrow G$;
2. Find a vertex v with indegree (the number of directed edges pointing to v) 0 in H , assign to it the number i . If no such vertex v exists, order the remaining vertices in H randomly and then STOP.
3. $H \leftarrow H - v, i \leftarrow i + 1$, if H is not empty, goto step 2.

Such an order $<$ is obstruction-free and thus is perfect. Therefore, G is perfectly orderable. $\square$

2.8.10 Intersection of Two Threshold Graphs

Definition 2.14 *A graph G is the intersection of two threshold graphs if there are two threshold graphs G_1 and G_2 such that $e \in E(G) \iff e \in E(G_1)$ and $e \in E(G_2)$.*

Hammer and Mahadev proved [10] that intersections of two threshold graphs are perfectly orderable.

2.8.11 D-Graphs

Definition 2.15 *A vertex x is a d -vertex if for any edge yz with $\{y, z\} \cap N(x) = \emptyset$, y and z are comparable. A graph G is a D -graph if each of its induced subgraphs contains a d -vertex.*

Hoàng ([12]) showed that the class of D -graphs contains all graphs with the Dilworth number at most three and all cotriangulated graphs. He also proved that every D -graph is perfectly orderable, and used this result to prove the following theorem:

Theorem 2.7 *Let G_1 be a threshold graph and let G_2 be a graph containing no induced P_4 and no induced C_4 . Then the union of G_1 and G_2 is perfectly orderable.*

□

A polynomial algorithm with the complexity of $O(nm)$ has been developed to recognize D -graphs ([12]).

2.8.12 Brittle Graphs

Definition 2.16 *Let $abcd$ be a P_4 of a graph G . The vertices a, d are called endpoints of the P_4 and the vertices b, c are called midpoints of the P_4 . G is called brittle if each induced subgraph H of G contains a vertex which is not an endpoint or a midpoint of a P_4 in H .*

Chvátal introduced brittle graphs and pointed out that they are perfectly orderable. For more information on brittle graphs, see ([14]). The class of brittle graphs

contains all P_4 -simplicial graphs, all P_4 -indifference graphs, all triangulated graphs and their complements, all Raspail graphs, all Welsh-Powell perfect graphs, all Matula perfect graphs and all graphs with Dilworth number at most three.

Chapter 3

INVESTIGATION OF THE PERFECT ORDERABILITY OF THE UNION OF TWO GRAPHS

With the motivation to generalize Theorem 2.6 (that the union of two threshold graphs is perfectly orderable), we study the perfect orderability of the unions of the two graphs, in which some or all of the three induced subgraphs P_4 , C_4 and $2K_2$ are forbidden. For three of the all 28 possible cases of the unions, we pose the conjecture that they form a new subclass of perfectly orderable graphs. Two theorems are proved to support our conjecture.

3.1 Motivation

In the last chapter, we introduced a subclass of perfectly orderable class, the union of the two threshold graphs, and also mentioned that by allowing a $2K_2$ in one of the graphs, Hoàng used the property of D -graph to prove the Theorem 2.7, which generalizes the Theorem 2.6.

We know that a threshold graph is a graph that contains no P_4 , $2K_2$ or C_4 as an induced subgraph. Given two graphs G_1 and G_2 , Theorem 2.7 implies that the

restriction that both G_1 and G_2 be threshold in Theorem 2.6 is stronger than necessary for the union $G = G_1 \cup G_2$ to be perfectly orderable. If we somehow weaken the restriction on G_1 and (or) G_2 , will the union G still be perfectly orderable? With this question in mind, we are going to list all the possible unions of G_1 and G_2 , in which we forbid 1 – 3 of these three structures (P_4 , $2K_2$ and C_4) as induced subgraphs, and study in each case the perfect orderability of G .

3.1.1 The List of Problems

We use Table 3.1 to list our problems:

3.1.2 Explanation of the List

Although there are 28 different cases of two graphs G_1 and G_2 in the list, many of them can be easily eliminated by finding a counter-example showing that the union G is not necessarily perfectly orderable.

• Counter-example 1:

Figure 3.1

The graph in Figure 3.1 is not perfectly orderable. (If we direct $a \rightarrow h$, then two edges gc and gf are forced to be directed from $g \rightarrow c$ and $g \rightarrow f$, resulting in an

Case NO.	G_1 contains no	G_2 contains no	$G_1 \cup G_2$ perfectly orderable?	Reason
1	$P_4, C_4, 2K_2$	P_4	NO	Figure 3.1
2	$P_4, C_4, 2K_2$	C_4	NO	C_5
3	$P_4, C_4, 2K_2$	$2K_2$	NO	C_5
4	$P_4, C_4, 2K_2$	P_4, C_4	YES	Theorem 2.7
5	$P_4, C_4, 2K_2$	$P_4, 2K_2$	?	
6	$P_4, C_4, 2K_2$	$C_4, 2K_2$	NO	C_5
7	$P_4, C_4, 2K_2$	$P_4, C_4, 2K_2$	YES	Theorem 2.6
8	P_4, C_4	P_4	NO	case 1
9	P_4, C_4	C_4	NO	case 2
10	P_4, C_4	$2K_2$	NO	case 3
11	P_4, C_4	P_4, C_4	NO	C_5
12	P_4, C_4	$P_4, 2K_2$	?	
13	P_4, C_4	$C_4, 2K_2$	NO	case 6
14	$P_4, 2K_2$	P_4	NO	case 1
15	$P_4, 2K_2$	C_4	NO	case 2
16	$P_4, 2K_2$	$2K_2$	NO	case 3
17	$P_4, 2K_2$	$P_4, 2K_2$	?	
18	$P_4, 2K_2$	$C_4, 2K_2$	NO	case 6
19	$C_4, 2K_2$	P_4	NO	case 1
20	$C_4, 2K_2$	C_4	NO	case 2
21	$C_4, 2K_2$	$2K_2$	NO	case 3
22	$C_4, 2K_2$	$C_4, 2K_2$	NO	case 6
23	P_4	P_4	NO	case 1
24	P_4	C_4	NO	case 2
25	P_4	$2K_2$	NO	case 3
26	C_4	C_4	NO	case 2
27	C_4	$2K_2$	NO	case 3
28	$2K_2$	$2K_2$	NO	case 3

Table 3.1: The List of the Union of Two Graphs.

impossible orientation on edge de without obstruction.)

In this Figure 3.1, we see that G_1 is a threshold graph, and G_2 is P_4 -free, so it is a counter-example to case 1 in the list. Since restrictions on G_1 are weaker in cases 8, 14, 19 and 23 than in case 1, we must say "NO" to all these cases.

• Counter-example 2 : C_5

Figure 3.2: C_5 .

C_5 is a well-known simple graph that is not perfectly orderable. If any of these unions in the list contains a C_5 as induced subgraph, then they are also not perfectly orderable.

In Figure 3.3, C_5 is written as the union of G_1 and G_2 such that G_1 is threshold and G_2 is $2K_2$ -free and C_4 -free. So it is a counter-example to cases 2, 3 and 6 in the list. It follows that the answers to cases 9, 10, 13, 15, 16, 18, 20-22, 24-28 are "NO".

In Figure 3.4, C_5 is written as the union of G_1 and G_2 such that G_1 and G_2 are both P_4 -free and C_4 -free. So it is a counter-example to cases 11 in the list.

Up to this point, we have given a definitive answer "YES" or "NO" to 25 out of 28 cases in the list, yet there are still 3 cases labeled with "?" that we cannot so easily answer.

Let us list again these three cases for clarity:

Figure 3.3: C_5 .

Figure 3.4: C_5 .

1. Case 5: G_1 is threshold, G_2 is P_4 -free and $2K_2$ -free;
2. Case 12: G_1 is P_4 -free and C_4 -free, G_2 is P_4 -free and $2K_2$ -free;
3. Case 17: G_1 and G_2 are both P_4 -free and $2K_2$ -free.

It is obvious that the union G in case 5 is a subclass of the unions in both cases 12 and 17.

3.2 The Perfect Orderability of the Union of Two Graphs in Three Cases

In this section, we shall investigate in more detail the perfect orderability of the union of two graphs in the cases 5, 12, 17. First, we verify that no known minimally non-perfectly orderable graphs can be written as the union of two graphs with the properties in these three cases. Second, we verify that no known class of perfectly orderable graphs can contain such unions. Third, we pose the conjecture that such three kinds of unions are new classes of perfectly orderable graphs, and with an extra restriction that G_1 and G_2 are edge disjoint, we prove the union in case 17. hence case 5, is perfectly orderable.

3.2.1 Some Minimally Non-perfectly Orderable Graphs

In this subsection, we introduce some known minimally non-perfectly orderable graphs and show that they cannot be the unions of two graphs with the properties in case 5, 12 and 17.

• Odd Hole

Definition 3.1 *An odd hole is a chordless cycle with odd length at least 5, namely C_{2k+1} ($k \geq 2$).*

C_5 is the simplest odd hole, which we used as a counter-example to many cases in the list, as explained in the last subsection. To show that the odd hole is not a counter-example to the three unanswered cases, we prove the following observation:

Observation 3.1 *Let G_1 be a graph containing no P_4 , no $2K_2$ as induced subgraphs, and G_2 be a graph containing no P_4 as an induced subgraph, then the union $G = G_1 \cup G_2$ contains neither C_k ($k \geq 5$) nor P_k ($k \geq 8$).*

Proof: Suppose that G contains a C_k ($k \geq 5$) or a P_k ($k \geq 8$). Since G_1 contains no P_4 and $2K_2$ as induced subgraphs, it is easy to see that G_1 can be of only two forms, P_2 and P_3 (Figure 3.5) in C_k ($k \geq 5$) or P_k ($k \geq 8$) (not considering the isolated vertices of G_1).

Figure 3.5: Possible G_1 .

But in either case, $C_k - G_1$ or $P_k - G_1$ contains a P_4 and is a subgraph of G_2 , a contradiction. $\square$

- **Anti-hole**

Recall that $\overline{G}$ is the complement of a graph G .

Definition 3.2 An anti-hole $\overline{C_k}$ is the complement of the cycle C_k .

Theorem 3.1 Anti-hole $\overline{C_k}$ ($k \geq 5$) is not perfectly orderable.

Proof: By contradiction. Suppose there is an admissible acyclic orientation on $\overline{C_k}$ ($k \geq 5$).

Let the vertices of an anti-hole $\overline{C_k}$ ($k \geq 5$) be denoted as $v_0, v_1, \dots, v_{k-1}$, such that $v_i v_{i+1}$ is not an edge (subscript is taken modulo k).

Observation 3.2 For every edge $v_i v_{i+2} \in \overline{C_k}$, if $v_i \rightarrow v_{i+2}$, then $v_{i-1} \rightarrow v_{i+1}$; if $v_{i+2} \rightarrow v_i$, then $v_{i+3} \rightarrow v_{i+1}$.

Proof: Note that both $v_i v_{i+2} v_{i-1} v_{i+1}$ and $v_{i+2} v_i v_{i+3} v_{i+1}$ are P_4 s in $\overline{C_k}$. By assumption, there is no obstruction in $\overline{C_k}$. So if $v_i \rightarrow v_{i+2}$, there is a forcing $v_{i-1} \rightarrow$

Figure 3.6

v_{i+1} in the P_4 $v_i v_{i+2} v_{i-1} v_{i+1}$; if $v_{i+2} \rightarrow v_i$, there is a forcing $v_{i+3} \rightarrow v_{i+1}$ in the P_4 $v_{i+2} v_i v_{i+3} v_{i+1}$ (See Figure 3.6.) $\square$

We continue the proof of the theorem. Without loss of generality, let us assume $v_0 v_2$ is directed from v_0 to v_2 .

Observation 3.2 implies that if k is even, $\overline{C_k}$ will have a directed cycle $v_0 v_2 v_4 \dots v_{k-2} v_0$; if k is odd, $\overline{C_k}$ will have a directed cycle $v_0 v_2 v_4 \dots v_{k-1} v_1 v_3 \dots v_{k-2} v_0$ (Figure 3.7), a contradiction.

Figure 3.7: An anti-hole.

To show that an anti-hole is also not a counter-example to cases 5, 12, and 17, we shall prove the following theorem:

Theorem 3.2 *Let G_1 be a graph containing no P_4 , no $2K_2$ as induced subgraphs, and G_2 be a graph containing no P_4 as induced subgraph, then the union $G = G_1 \cup G_2$*

contains no $\overline{C_k}$ ($k \geq 5$).

The proof of Theorem 3.2 will be presented in section 3.3 of this chapter.

• Other Examples

Here, we give some more specific examples of minimally non-perfectly orderable graphs:

i)

Figure 3.8: Example 1.

In Figure 3.8, without loss of generality, we may assume that $6 \rightarrow 7$. There is a chain reaction forcing $8 \rightarrow 4$, $3 \rightarrow 9$, $10 \rightarrow 1$ and $2 \rightarrow 3$, whereupon the vertices 2, 3, 4 and 8 constitute an obstruction.

It is clear that the graph G in Figure 3.8 contains a P_8 , namely 678439(10)1. and using the Observation 3.1, we know that G cannot be the union of two graphs in case 5, 12 and 17.

ii)

The graph G in Figure 3.9 is the union of a $\overline{P_7}$ 1234567 and and a P_4 3895. Without loss of generality, we may assume $8 \rightarrow 9$. Then there is a chain reaction forcing $5 \rightarrow 7$, $4 \rightarrow 6$, $3 \rightarrow 5$, $2 \rightarrow 4$, $1 \rightarrow 3$ and $5 \rightarrow 1$ (by $8 \rightarrow 9$), whereupon there is a directed cycle 1351 in G .

Figure 3.9: Example 2.

To show that G cannot be the union of G_1 and G_2 in cases 5, 12 and 17, we shall prove the following two observations:

Observation 3.3 G cannot be the union of two graphs G_1 and G_2 in case 17.

Proof: By contradiction. Suppose G is the union of G_1 and G_2 in case 17, that is G_1 and G_2 are both P_4 -free and $2K_2$ -free.

We notice that there is a P_6 , namely 895746, in G . Since G_1 and G_2 are both P_4 -free and $2K_2$ -free, they can be of only two forms, P_2 or P_3 , in P_6 (not considering the isolated vertices). In either case, G_2 (or G_1) = $G - G_1$ (or G_2) contains a P_4 , a contradiction. $\square$

Observation 3.4 G cannot be the union of G_1 and G_2 in case 12.

Proof: By contradiction. Suppose G_1 and G_2 are in case 12, that is G_1 is P_4 -free and C_4 -free, G_2 is P_4 -free and $2K_2$ -free.

Consider the P_6 895746 induced in G . We must have two outside edges 89 and 46 belonging to G_1 only and the middle edge 57 belonging to G_2 only. (We say that an edge e belongs to G_i only if $e \in E(G_i) - E(G_j)$ with $i \neq j$.) In the C_4 8953, since 89 is in G_1 and G_1 contains no P_4 and C_4 , we can see that either i) 35, 38 $\in G_2$ only, and 59 $\in G_1$ only, or ii) 35, 95 $\in G_2$ only, and 38 $\in G_1$ only. Without loss of generality,

we may assume the first case occurs. Now, if $25 \in G_1$, then there is a P_4 2598 in G_1 ; if $25 \in G_2$, then there is a P_4 2538 in G_2 , a contradiction. $\square$

Since the classes of unions described in cases 12 and 17 contain all unions in case 5, the above two observations are sufficient to show G cannot be the union of two graphs in cases 5.

iii)

Figure 3.10: Example 3.

The graph G in Figure 3.10 is a minimally non-perfectly orderable graph (the forcing on the edges of G creates a directed cycle 1471).

Suppose $G = G_1 \cup G_2$ as in any of cases 5, 12, or 17. If both edges 34 and 45 belong to G_1 , then consider the P_5 23456 of G . Since G_1 contains no P_4 , the edges 23 and 56 must belong to G_2 and they form a $2K_2$ in G_2 , a contradiction. Otherwise, at least one of the edges 34 or 45 must be from G_2 , so by symmetry assume 34 is in G_2 . Similarly, at least one of the edges 67, or 78 is in G_2 . In each case, the set $\{3, 4, 6, 7\}$ (or $\{3, 4, 7, 8\}$) induces a $2K_2$ or P_4 in G_2 , a contradiction.

3.2.2 Comparison with Known Classes of Perfectly Orderable Graphs

In this subsection, we will show that the union of two graphs in cases 5, 12 and 17 does not belong to any known class of perfectly orderable graphs introduced in Chapter 1.

Since the union in case 5 has more strict restrictions on G_1 , it is sufficient to show that such a union does not belong to any class of perfectly orderable graphs. Next, we are going to give some examples to illustrate our conclusion.

- **Example 1**

Figure 3.11: Example 1.

In figure 3.11, the graph G is the union in case 5 (G_1 consists of the solid lines, G_2 consists of the dashed lines).

i) G is not a comparability graph.

It cannot be directed without constructing a semi-transitive orientation in Figure 2.5. (If we direct $b \rightarrow c$, then $d \rightarrow e$. Comparability orientations imply $f \rightarrow e$, $a \rightarrow e$ and $a \rightarrow b$, then we have a directed P_3 $a \rightarrow b, b \rightarrow c$ that is not a comparability orientation.)

ii) G is not a triangulated graph.

It contains a C_4 .

iii) G is not a co-triangulated graph.

The two edges ef and bc form a $2K_2$ in G , so there is a C_4 $becf$ in $\overline{G}$, which is not a triangulated graph.

iv) G is not a P_4 -comparability graph.

Suppose that G is a P_4 -comparability graph. Without loss of generality, we may assume $b \rightarrow c$. This implies $d \rightarrow e$ and $e \rightarrow a$. But then one of the P_4 's $feab$ or $fedc$ is not of Type 3, a contradiction.

v) G is not a P_4 -indifference graph.

In Figure 3.11, if P_4 $eabc$ is of Type 2, P_4 $bcde$ cannot be of Type 2, otherwise there will be a directed cycle $bcdeab$.

vi) G is not a Raspail graph.

G contains a *a house* which is a forbidden subgraph in Raspail graphs.

vii) G is not a Welsh-Powell perfect nor a Matula perfect graph.

A house is also a forbidden structure in both Welsh-Powell and Matula perfect graphs.

viii) G is not a union of two threshold graphs.

This is obvious.

• Example 2

The graph G in Figure 3.12 (a) is the union of two threshold graphs G_1 and G_2 , with G_1 consisting of the edges ab, ac, bc and G_2 consisting of the remaining edges.

Claim 3.1 G is not an intersection of two threshold graphs.

Proof: Suppose G is the intersection of two threshold graphs G_1, G_2 .

Observation 3.5 Let e_1, e_2 be two edges that induce a $2K_2$ in G . Then some edge e_i ($i = 1, 2$) must have one of its endpoints being adjacent to the two endpoints of e_j

Figure 3.12

($j \neq i$) in G_1 and the other endpoint being adjacent to the two endpoints of e_j in G_2 (Figure 3.13). $\square$

(In Figures 3.13 and 3.12 (b), the thick lines denote edges in both G_1 and G_2 , the solid lines denote edges in G_1 , and the dashed lines denote edges in G_2 .)

Figure 3.13

Consider the $2K_2$ formed by $\{a', c', a, c\}$. By Observation 3.5, we may assume that $aa', ac' \in E(G_1)$ and $ca', cc' \in E(G_2)$. Now consider the $2K_2$ formed by $\{a, b, a', c'\}$, Observation 3.5 implies that $ba', bc' \in E(G_2)$. But the set $\{a', c', b, c\}$ induces a $2K_2$ in G_1 , a contradiction. $\square$

We conclude that there exists a union in cases 5, 12, and 17 that does not belong to the class of intersections of two threshold graphs.

• Example 3

Figure 3.14: A non-D-graph.

The graph G in Figure 3.14 is a union of G_1 (in solid line) and G_2 (in dashed line). It is easy to verify that G_1 is threshold, and G_2 is P_4 -free and $2K_2$ -free.

The graph G

i) has Dilworth number greater than 3.

Let $S = \{5, 6, 3, 1\}$. It is easy to verify that the vertices in S are pairwise incomparable, so the Dilworth number of G is at least 4.

ii) is not a D -graph.

Recall that a d -vertex w is such a vertex that for every edge ab with $\{a, b\} \cap N(w) = \emptyset$, a and b are comparable. There is no d -vertex in G (for vertices 1, 2, 3 and 4, we have that 5 and 6 are incomparable; for vertices 3' and 4', we have that 5' and 6' are incomparable; for vertices 5, 6, 7 and 8 (respectively 5', 6', 7', and 8'), we have that 3 (3') and 1 are incomparable).

• Example 4

Figure 3.15

The graph G shown in Figure 3.15 is a union of two threshold graphs. The reader may easily check that each vertex is the endpoint of some P_4 and the midpoint of some other P_4 . Thus G is not brittle (and therefore not P_4 -simplicial).

The above four examples show that no known class of perfectly orderable graphs contains all unions of two graphs in case 5, or 12, or 17.

3.2.3 Conjectures

Since we are unable to find a non perfectly orderable graph that can be written as the union of two graphs in case 5, or 12, or 17, we would like to propose the following conjecture:

Conjecture 1 *If G is the union of two graphs G_1 and G_2 satisfying any of the following conditions:*

1. G_1 is threshold, G_2 is P_4 -free and $2K_2$ -free;
2. G_1 is P_4 -free and C_4 -free, G_2 is P_4 -free and $2K_2$ -free;

3. G_1 and G_2 are both P_4 -free and $2K_2$ -free.

then G is perfectly orderable.

If Conjecture 1 is true, then the three classes of graphs described by it would form new classes of perfectly orderable graphs (by the results in subsection 3.2.2).

Although we have not found a way to prove this conjecture, we do prove, jointly with Hoàng, the following theorem by adding the extra constraint that G_1 and G_2 are edge-disjoint for case 3.

Theorem 3.3 *If G_1 and G_2 are both P_4 -free and $2K_2$ -free, and G_1 and G_2 are edge disjoint, then $G = G_1 \cup G_2$ is perfectly orderable.*

The proof of this theorem is given in the next section.

3.3 Proof of the Theorems

Theorem 3.4 *Let G_1 be a graph containing no P_4 , no $2K_2$ as induced subgraphs, and G_2 be a graph containing no P_4 as induced subgraph, then $G = G_1 \cup G_2$ contains no $\overline{C}_k$ ($k \geq 5$).*

Proof: By contradiction. Suppose G contains $\overline{C}_k$ ($k \geq 5$). Number the vertices of $\overline{C}_k$ as $v_0, v_1 \dots v_{k-1}$, such that $v_i v_{i+1}$ is not an edge (subscript is taken modulo k). Since it is easy to verify that the Theorem is true for $k = 5$, we may assume that $k > 5$.

Before making any further argument, we are going to introduce the following observation, which is frequently used in the proof (whenever we say an edge e is in G_l ($l = 1, 2$) only, we mean $e \in E(G_l) - E(G_t)$, $t \neq l$):

Observation 3.6 *In any induced C_4 of $\overline{C}_k$ ($k \geq 5$), namely $abcd$, if ab is in G_1 only, and bc is in G_2 only, then ad is in G_1 only, and cd is in G_2 only (Figure 3.16). $\square$*

Figure 3.16

Claim 3.2 For any four consecutive vertices $v_i, v_{i+1}, v_{i+2}, v_{i+3}$ ($i = 0, 1, \dots, k-1$), the two edges $v_i v_{i+2}$ and $v_{i+1} v_{i+3}$ cannot be in the same G_l ($l = 1, 2$).

Proof: Suppose $v_i v_{i+2}$ and $v_{i+1} v_{i+3}$ are in the same G_l . we need to consider only four cases:

1. They are both in G_1 only;
2. They are both in G_2 only;
3. One is in both G_1 and G_2 , the other one is in G_l ($l = 1, 2$) only;
4. They are both in G_1 and G_2 .

Since G_1 contains no P_4 and $2K_2$, it is obvious that case 1 and case 4 are not possible. We are now going to show that case 2 and case 3 are not possible either.

case 2: Suppose $v_i v_{i+2}$ and $v_{i+1} v_{i+3}$ are in G_2 only. Since $v_{i+1} v_{i+3} v_i v_{i+2}$ is a P_4 of G , $v_i v_{i+3}$ must be in G_1 only. Consider the C_4 $v_{i-1} v_{i+2} v_i v_{i+3}$: using Observation 3.6, we have $v_{i-1} v_{i+2}$ belonging to G_2 only and $v_{i+3} v_{i-1}$ belonging to G_1 only. Now $v_{i-1} v_{i+1}$ cannot belong to either G_1 or G_2 (if $v_{i-1} v_{i+1}$ is in G_1 then $v_{i+1} v_{i-1} v_{i+3} v_i$ is a P_4 in G_1 ; if $v_{i-1} v_{i+1}$ is in G_2 then $v_{i+1} v_{i-1} v_{i+2} v_i$ is a P_4 in G_2), a contradiction (see Figure 3.17.)

case 3: Suppose $v_i v_{i+2}$ is in both G_1 and G_2 , $v_{i+1} v_{i+3}$ is in G_l ($l = 1, 2$) only.

Figure 3.17

Since G_1 contains no P_4 and $2K_2$, $v_{i+1}v_{i+3}$ cannot be in G_1 (otherwise, the set $\{v_i, v_{i+1}, v_{i+2}, v_{i+3}\}$ induces a P_4 or $2K_2$ in G_1), and must be in G_2 only, and $v_i v_{i+3}$ is in G_1 only.

Figure 3.18

Consider the C_4 $v_i v_{i+3} v_{i+1} v_{i+4}$: Observation 3.6 implies that $v_{i+4} v_{i+1}$ is in G_2 only, and $v_i v_{i+4}$ is in G_1 only. For $j = i + 4, i + 5, \dots, k - 1, 0, 1, \dots, i - 3$, by considering the C_4 $v_i v_j v_{i+1} v_{j+1}$, we see that every edge $v_i v_{j+1}$ is in G_1 only and every edge $v_{i+1} v_{j+1}$ is in G_2 only. Consider the P_4 $v_i v_{i-2} v_{i+1} v_{i-1}$: with $v_{i+1} v_{i-2}$ belonging to G_2 only and

$v_{i-2}v_i$ belonging to G_1 only, it is obvious that $v_{i+1}v_{i-1}$ is in G_2 only. So all the edges going out from v_i (except for v_iv_{i+2}) are in G_1 only, and all the edges going out from v_{i+1} are in G_2 only (see Figure 3.18.)

Figure 3.19

Consider the $P_4 v_{i+3}v_{i+1}v_{i+4}v_{i+2}$: with both edges $v_{i+1}v_{i+3}$ and $v_{i+1}v_{i+4}$ belonging to G_2 only, it is clear that $v_{i+2}v_{i+4}$ is in G_1 only. For $j = i+4, \dots, k-1, 0, 1, \dots, i-2$, consider the $C_4 v_{i+1}v_jv_{i+2}v_{j+1}$, where $v_{i+1}v_j$ and $v_{i+1}v_{j+1}$ are in G_2 only and $v_{i+2}v_j$ is in G_1 only, we have that every edge $v_{i+2}v_{j+1}$ is in G_1 only. For $j = i, i-1, \dots, 0, k-1, \dots, i+4$, by considering the $C_4 v_{i+3}v_jv_{i+2}v_{j-1}$, where $v_{i+3}v_j$, v_jv_{i+2} and $v_{i+2}v_{j-1}$ are in G_1 , we see that $v_{i+3}v_{j-1}$ is in G_1 . Now consider the $P_4 v_{i+3}v_{i+5}v_{i+2}v_{i+4}$: we see that all three edges belong to G_1 , a contradiction. This completes the proof of Claim 3.2 (see Figure 3.19.)

Observation 3.7 In any induced $C_4 abcd$ of $\overline{C_k}$, if ab is in G_1 only, and ad and bc are in G_2 , then ac and bd must belong to G_1 .

Proof: It is clear that if ad and bc are both in G_2 only, we either have a $2K_2$ ab and cd in G_1 , or a $P_4 adcb$ in G_2 , a contradiction. Without loss of generality, assume ad belongs to G_1 and bc does not. Then either $adcb$ is a P_4 in G_2 or $badc$ is a P_4 in G_1 . $\square$

Figure 3.20

Now, we continue the proof of the theorem.

By Claim 3.2 we may assume, without loss of generality, that v_0v_2 is in G_1 only (if v_0v_2 is in G_2 then v_1v_3 is in G_1 only, by renumbering the vertices of $\overline{C_k}$, we could arrive at the same conclusion). Again, Claim 3.2 implies that v_1v_{k-1} and v_1v_3 are in G_2 only. Consider the C_4 $v_0v_2v_{k-1}v_3$ in G . By Observation 3.7, one of the following two cases must occur: (i) either v_0v_3 belongs to G_1 only or v_2v_{k-1} belongs to G_1 only, or (ii) both v_0v_3 and v_2v_{k-1} belong to G_1 and G_2 . Suppose that case (ii) occurs. Then we have v_3v_{k-1} belonging to G_1 only; for otherwise $v_0v_3v_{k-1}v_2$ is a P_4 in G_2 . But now $v_3v_1v_{k-1}v_2$ is a P_4 in G_2 . Thus we know that case (i) must occur. Without loss of generality, we may assume that v_0v_3 is in G_1 only (see Figure 3.20).

For $j = 3, 4, \dots, k-3$, consider the C_4 $v_0v_jv_1v_{j+1}$: with v_0v_j belonging to G_1 only and v_1v_j belonging to G_2 only, by Observation 3.6 we have v_0v_{j+1} belonging to G_1 only and v_1v_{j+1} belonging to G_2 only (see Figure 3.21.)

Consider the P_4 $v_3v_1v_4v_2$: with two edges v_3v_1 and v_1v_4 belonging to G_2 only, we must have v_2v_4 belonging to G_1 only. For $j = 4, 5, \dots, k-2$, by considering the C_4 $v_1v_jv_2v_{j+1}$, with both edges v_jv_1 and v_1v_{j+1} belonging to G_2 only, and v_2v_j belonging to G_1 only, we have v_2v_{j+1} belonging to G_1 only (see Figure 3.22.)

Figure 3.21

Consider the C_4 $v_0v_2v_{k-1}v_3$: with v_3v_0 , v_0v_2 , and v_2v_{k-1} belonging to G_1 , we have v_3v_{k-1} belonging to G_1 . For $j = k-2, k-3, \dots, 5$, considering the C_4 $v_2v_jv_3v_{j-1}$: with three edges v_3v_{j-1} , $v_{j-1}v_2$ and v_2v_j belonging to G_1 , we must have v_3v_j belonging to G_1 as well. Now, look at the P_4 $v_3v_5v_2v_4$, all three edges of P_4 are in G_1 , a contradiction to the definition of G_1 (see Figure 3.23.) $\square$

Before we start the proof of Theorem 3.3, we restate it:

Theorem 3.3 *If G_1 and G_2 are both P_4 -free and $2K_2$ -free, and G_1 and G_2 are edge disjoint, then $G = G_1 \cup G_2$ is perfectly orderable.*

Proof: Let G_1 , G_2 and G be as specified in the statement of the Theorem.

Claim 3.3 *For every edge $ab \in E(G_1)$ (respectively $ab \in E(G_2)$), the two vertices a , b must be comparable in G_2 (respectively G_1).*

Proof: Suppose $ab \in E(G_1)$ and a, b are not comparable in G_2 . It follows that there exist two edges $ad \in E(G_2)$, $bc \in E(G_2)$, such that $ac \notin E(G_2)$ and $bd \notin E(G_2)$. Then there will be either a $2K_2$ or P_4 in G_2 , which is a contradiction (see Figure 3.24.)

$\square$

Figure 3.22

Now define a partial orientation on G (we say that a strictly dominates b if a dominates b but b does not dominate a).

1. $\forall ab \in E(G_1)$, direct $a \rightarrow b$ if and only if a strictly dominates b in G_2 ;
2. $\forall cd \in E(G_2)$, direct $c \rightarrow d$ if and only if c strictly dominates d in G_1 .

Claim 3.4 G has no obstruction under the partial orientation.

Proof: Since G_1 and G_2 are P_4 -free and $2K_2$ -free, the P_4 $abcd$ in G , if there is any, must be of the following two forms (see Figure 3.25):

1. $ab, bc \in E(G_1), cd \in E(G_2)$,
2. $ab \in E(G_1), bc, cd \in E(G_2)$.

By definition of the partial orientation, we have $c \rightarrow d$ in 1) and $b \rightarrow a$ in 2). In either case, there is no obstruction. $\square$

Claim 3.5 The partial orientation creates no cycle in G .

Figure 3.23

Figure 3.24

Proof: Clearly, there are no cycles in the same G_l ($l = 1, 2$) since the domination relation is transitive.

Now we are going to show there is no *mixed* cycle, the cycle made up of directed edges in G_1 and G_2 .

Observation 3.8 If $\overrightarrow{ab} \in E(G_1)$ (respectively, $\overleftarrow{ab} \in E(G_2)$) and $\overrightarrow{bc} \in E(G_2)$ (respectively, $\overleftarrow{bc} \in E(G_1)$), then there must be $\overrightarrow{ac} \in E(G_2)$ (respectively, $\overleftarrow{ac} \in E(G_1)$).

Proof: $\overrightarrow{ab} \in E(G_1)$ implies a strictly dominates b in G_2 . Since $\overrightarrow{bc} \in E(G_2)$, we have $ac \in E(G_2)$, and by Claim 3.3, a and c are comparable in G_1 . Since $ab \in E(G_1)$ and $cb \notin E(G_1)$, a strictly dominates c and there is an orientation from a to c (see Figure 3.26.) $\square$

Figure 3.25: Two possible P_4 .

Figure 3.26

Observation 3.8 implies that there is no mixed cycle on three vertices.

Suppose there is a cycle in G , then it must be a mixed cycle. Let C be a cycle $v_0v_1\dots v_{l-1}$ with smallest length l . Select v_i ($i \in \{0, 1, \dots, l-1\}$), such that $v_{i-1}\overline{v_i} \in E(G_1)$, and $v_i\overline{v_{i+1}} \in E(G_2)$. By the Observation 3.8, we have $v_{i-1}\overline{v_{i+1}} \in E(G_2)$. Then there is a new cycle $v_0v_1\dots v_{i-1}v_{i+1}\dots v_{l-1}$ of length $l-1$, contradicting with the assumption that C is the smallest cycle. $\square$

Based on this acyclic partial orientation, we can easily construct a linear order $<$ on G by the following procedure:

1. $i \leftarrow 1, H \leftarrow G$;
2. Find a vertex v in H with indegree 0 and assign the order i to v ;
if no such a v exists, order the remaining vertices in H randomly and then exit;
3. $i \leftarrow i + 1, H \leftarrow H - v$, if $H \neq \emptyset$ goto step 2.

Such an order $<$ is sure to be obstruction-free, therefore, G is perfectly orderable.

□

3.4 Discussion of the Proof of Theorem 3.3

In the proof of Theorem 3.3, we first impose a partial orientation on the edges of G according to a certain rule, which guarantees the orientation to be unique on each edge, obstruction-free and acyclic, later construct a perfect linear order $<$ on the vertices of G based on this partial orientation, thus get the conclusion that G is perfectly orderable. The key point is how to find such an effective rule for orienting edges. We found it because we noticed the fact that in the two possible P_4 s in Figure 3.25, if we direct the edges in G_1 (respectively G_2) according to the strict domination in G_2 (respectively G_1), it guarantees there is at least one wing of P_4 going out from the joint to the tip, preventing an obstruction. With the conditions provided in Theorem 3.3, we are able to show that such a rule is good in the since it can produce a perfect order on G . Here, the constraint that G_1 and G_2 are edge-disjoint is important, otherwise such a rule will not work, even for the union of two threshold graphs. In the graph of Figure 3.27, a union of two threshold graphs (one in solid lines and one in dashed lines), this strict domination rule causes a directed cycle.

Realizing the effect of such a constraint on the rule, we naturally ask this question: can a union G of two threshold graphs G_1 and G_2 be decomposed into two edge-disjoint graphs G'_1 and G'_2 , such that G'_1 and G'_2 are P_4 -free and $2K_2$ -free? Unfortunately, a counter-example is found in Figure 3.28 to give a "NO" answer to the question above.

The graph in Figure 3.28 is a union of two threshold graphs with a common edge in both G_1 and G_2 . Since both G'_1 and G'_2 are P_4 -free and $2K_2$ -free, if we put edge ef in G'_1 , all the edges in G_2 except ab must be put in G'_2 ; similarly, if $e'f'$ is in G'_2 ,

Figure 3.27: A counter-example to the rule.

all the edges in G_1 except ab must be in G'_1 . If ab is not in G'_2 , there will be a P_4 $ac'bd'$ in G'_2 ; if ab is not in G'_1 , there will be a P_4 $dacb$ in G'_1 , a contradiction.

Figure 3.28: A counter-example to the question.

Bibliography

- [1] K. Appel and W. Haken. Every planar map is 4-colorable-1: Discharging. *Ill. J. Math.*, 21:429–490, 1977.
- [2] K. Appel and W. Haken. Every planar map is 4-colorable-2: Reducibility. *Ill. J. Math.*, 21:491–567, 1977.
- [3] C. Berge and V. Chvátal, editors. *Topics on perfect graphs*. North-Holland, Amsterdam, 1984.
- [4] V. Chvátal and P.L. Hammer. Aggregation of inequality in integer programming. *Annals of Disc. Math.*, 1:145–162, 1977.
- [5] V. Chvátal, C.T. Hoàng, N.V.R. Mahadev, and D. de Werra. Four classes of perfectly orderable graphs. *J. Graph Theory*, 11(4):481–495, 1987.
- [6] D.J.A. Welsh and M.B. Powell. An upper bound on the chromatic number of a graph and its application to timetabling problems. *Computer J.*, 10:85–87, 1967.
- [7] M. Garey and D.S. Johnson. *Computer and Intractability : A Guide to the Theory of NP-completeness*. W.H. Freeman, San Francisco, 1979.
- [8] A. Ghouila-Houri. Caractérisation des graphes non orientés dont on peut orienter les arêtes de manière à obtenir le graphe d'une relation d'ordre. *C.R. Acad. Sci. Paris*, 254:1370–1371, 1962.
- [9] M.C. Golumbic. *Algorithmic graph theory and perfect graphs*. Academic Press, New York, 1980.
- [10] P. L. Hammer and N. V. R. Mahadev. Bithreshold graphs. *SIAM J. Appl. Math.*, pages 497–506, 1985.
- [11] A. Hertz and D. de Werra. Les graphes bipolarisables. Report ORWP, Swiss Federal Institute of Technology in Lausanne, 1986.
- [12] C.T. Hoàng. A note on perfectly orderable graphs. *Discrete Applied Math.*, 1996.

- [13] C.T. Hoàng. On the complexity of recognizing a class of perfectly orderable graphs. *Discret Applied Math.*, pages 219–226, 1996.
- [14] C.T. Hoàng and N. Khouzam. On brittle graphs. *J. Graph Theory*, 12(3):391–404, 1988.
- [15] C.T. Hoàng and B.A. Reed. Some classes of perfectly orderable graphs. *J. Graph Theory*. 13(4):445–463, 1989.
- [16] C. Lund and M. Yannakakis. On the hardness of approximating minimization problems. In *Symposium on the Theory of Computing*, pages 286–293. 1993.
- [17] M. Middendorf and F. Pfeiffer. On the complexity of recognizing perfectly orderable graphs. *Discrete Math.*, 80(3):327–333, 1990.
- [18] Brooks R.L. On coloring the nodes of a network. *Proc. Cambridge Phil. Soc.*, pages 194–197. 1941.
- [19] Maciej M Syslo, Narsingh Deo, and Janusz S. Kowalik. *Discrete Optimization Algorithms*. Prentice-Hall, Inc., Englewood Cliffs, New Jersey 07632, 1983.

IMAGE EVALUATION TEST TARGET (QA-3)

APPLIED IMAGE, Inc
1653 East Main Street
Rochester, NY 14609 USA
Phone: 716/482-0300
Fax: 716/288-5989

© 1993, Applied Image, Inc., All Rights Reserved